

Convex isoperimetric sets in the Heisenberg group

Roberto Monti*

Dipartimento di Matematica Pura ed Applicata,
Università di Padova, Via Belzoni 7,
35100 Padova, Italy.
email: monti@math.unipd.it

Matthieu Rickly†

Institute of Mathematics,
University of Bern, Sidlerstrasse 5,
3012 Bern, Switzerland.
email: matthieu.rickly@math.unibe.ch

July 26, 2006

Abstract

We characterize convex isoperimetric sets in the Heisenberg group endowed with horizontal perimeter. We first prove Sobolev regularity for a certain class of $\mathbb{R}^2$ -valued vector fields of bounded variation in the plane related to the curvature equations. Then, by an approximation-reparameterization argument, we show that the boundary of convex isoperimetric sets is foliated by geodesics of the Carnot-Carathéodory distance.

1 Introduction

We identify the Heisenberg group $\mathbb{H}$ with $\mathbb{R}^3 = \mathbb{C} \times \mathbb{R}$ endowed with the group law

$$(z, t)(z', t') = (z + z', t + t' + 2\operatorname{Im}z\bar{z}'), \quad (1.1)$$

where $t, t' \in \mathbb{R}$, $z = x + iy$, $z' = x' + iy' \in \mathbb{C}$ and $\operatorname{Im}z\bar{z}' = yx' - xy'$. The Lie algebra of left-invariant vector fields is spanned by

$$X = \frac{\partial}{\partial x} + 2y \frac{\partial}{\partial t}, \quad Y = \frac{\partial}{\partial y} - 2x \frac{\partial}{\partial t} \quad \text{and} \quad T = \frac{\partial}{\partial t}, \quad (1.2)$$

*The first author would like to thank the Institute of Mathematics of the University of Bern, where part of the work was done.

†The second author would like to thank the Institute for Mathematical Research of the ETH Zürich, the Institute of Mathematics of the University of Bern and the Department for Pure and Applied Mathematics of the University of Padova.

and the distribution spanned by X and Y , called horizontal distribution, generates the Lie algebra by brackets.

The natural volume in $\mathbb{H}$ is given by the Haar measure, which, up to a positive factor, coincides with Lebesgue measure $\mathcal{L}^3$ in $\mathbb{R}^3$. Lebesgue measure can also be recovered as the Riemannian volume induced by the left-invariant metric for which X, Y and T are orthonormal. We denote the volume of a measurable set $A \subseteq \mathbb{H}$ by $\mathcal{V}(A) = \mathcal{L}^3(A)$.

The horizontal perimeter (or simply perimeter) of a measurable set $A \subseteq \mathbb{H}$ is

$$\mathcal{P}(A) = \sup \left\{ \int_A (X\varphi_1 + Y\varphi_2) d\mathcal{L}^3 \mid \varphi_1, \varphi_2 \in C_c^1(\mathbb{R}^3), \varphi_1^2 + \varphi_2^2 \leq 1 \right\}. \quad (1.3)$$

If $\mathcal{P}(A) < +\infty$, the set A is said to be of finite perimeter. Perimeter is left-invariant and 3-homogeneous with respect to the group of dilations $\delta_\lambda : \mathbb{H} \rightarrow \mathbb{H}$ given by

$$\delta_\lambda(z, t) = (\lambda z, \lambda^2 t), \quad \lambda > 0, \quad (1.4)$$

that is $\mathcal{P}(\delta_\lambda(A)) = \lambda^3 \mathcal{P}(A)$. Definition (1.3) of perimeter, which is modelled on De Giorgi's notion of perimeter in Euclidean spaces, was introduced in [GN] (see also [FGW]). If A is smooth (e.g. of class C^2), the perimeter coincides with the Minkowski content and with the 3-dimensional Hausdorff measure, precisely

$$\mathcal{P}(A) = \lim_{\varepsilon \downarrow 0} \frac{\mathcal{V}(A_\varepsilon) - \mathcal{V}(A)}{\varepsilon} = \mathcal{S}^3(\partial A). \quad (1.5)$$

Here, A_ε is the ε -neighbourhood of A with respect to Carnot-Carathéodory distance (we recall the definition in Section 2), and $\mathcal{S}^3$ is the spherical 3-Hausdorff measure, properly normalized, defined by means of the same distance. The equality of perimeter and Minkowski content is proved in [MSC], the equality of perimeter and $\mathcal{S}^3$ is proved in [FSSC]. Perimeter also admits an integral-geometric representation formula (see [M]).

Volume and perimeter are related via the isoperimetric inequality

$$\mathcal{V}(A) \leq c \mathcal{P}(A)^{4/3}, \quad (1.6)$$

where $c > 0$ is a constant and $A \subseteq \mathbb{H}$ is any measurable set with finite perimeter and volume. This inequality is proved by P. Pansu in [P1] and [P2] for smooth domains, with $\mathcal{S}^3(\partial A)$ instead of $\mathcal{P}(A)$. A set $A \subseteq \mathbb{H}$ with $0 < \mathcal{V}(A) < +\infty$ is called an isoperimetric set if it minimizes the isoperimetric ratio

$$\mathcal{I}(A) = \frac{\mathcal{P}(A)^{4/3}}{\mathcal{V}(A)}. \quad (1.7)$$

The existence of isoperimetric sets is proved by Leonardi and Rigot in [LR] by a concentration-compactness argument. Pansu notes that the boundary of a smooth isoperimetric set has “constant mean curvature” and that a smooth surface has “constant mean curvature” if and only if it is foliated by horizontal lifts of plane circles with constant radius. Then he conjectures that an isoperimetric set is obtained by

rotating around the center of the group a geodesic joining two points in the center. Recently, Pansu's conjecture reappeared in [LM].

Up to a left translation, the conjectured isoperimetric sets are rotationally symmetric, i.e. they are invariant with respect to isometries which leave the center invariant. However, the non commutative group law makes it difficult to prove by rearrangement or symmetrization that the isoperimetric ratio (1.7) is minimized by rotationally symmetric sets.

Assuming rotational symmetry, it is easy to determine the isoperimetric profile, as shown in [Mo] and [DGN1] in some special cases, and in [RR1] in the general case. In fact, it suffices to assume the rotational symmetry of a certain horizontal section (see [DGN2] and [R]). Isoperimetric sets can be also determined assuming regularity instead of symmetry. This is an important new result due to M. Ritoré and C. Rosales. In [RR2], they prove Pansu's conjecture for isoperimetric sets of class C^2 , without any symmetry assumption. However, a regularity theory for isoperimetric sets in the Heisenberg group does not yet exist (but see [CHMY] and [Pa2]).

It is worth mentioning that a 2-dimensional version of the problem is formulated and solved by the first author and Morbidelli in [MM]. In this setting, isoperimetric sets coincide with the section of the set (1.8) below with the $y = 0$ plane, properly scaled and translated. Finally, we refer the reader to the monograph [CDST] for a more detailed introduction to the isoperimetric problem in the Heisenberg group.

In this article, we characterize isoperimetric sets in the class of convex sets. A set $C \subseteq \mathbb{H}$ is convex if it is convex with respect to the usual vector space structure of $\mathbb{R}^3$. We do not assume any symmetry or regularity besides convexity.

Theorem 1.1 (Convex isoperimetric sets). *Up to a left translation and a dilation, any open isoperimetric set in $\mathbb{H}$ coincides with the set*

$$\left\{ (z, t) \in \mathbb{H} \mid |t| < \arccos |z| + |z| \sqrt{1 - |z|^2}, |z| < 1 \right\}. \quad (1.8)$$

The set in (1.8) is foliated by Heisenberg geodesics (see Section 2), it is globally of class C^2 , but it fails to be of class C^3 at the north and south poles $(0, \pm\pi/2)$. At these points, the plane spanned by the vector fields X and Y , the horizontal plane, is tangent to the boundary of the set.

In the first part of the paper, we describe the characteristic set of a convex set $C \subseteq \mathbb{H}$. We say that a point $p \in \partial C$ is characteristic if the horizontal plane at p is a supporting plane for C at p and we write $p \in \Sigma(C)$. We show that $\Sigma(C) = \Sigma^- \cup \Sigma^+$, where Σ^- , Σ^+ are two closed, disjoint, horizontal segments (possibly points). This is Theorem 3.5 in Section 3.

In Section 4, we derive the curvature equations by a variational argument. We consider a convex set

$$C = \{ (z, t) \in \mathbb{H} \mid z \in D, f(z) \leq t \leq g(z) \}, \quad (1.9)$$

where $D \subseteq \mathbb{R}^2$ is a closed convex set in the plane with nonempty interior, and $-g, f : D \rightarrow \mathbb{R}$ are convex functions. The intersection of $\Sigma(C)$ with the graph of f is a union of two (possibly empty) line segments. We denote its projection onto the xy -plane by $\Sigma(f)$.

If $C \subseteq \mathbb{H}$ is isoperimetric, then the function $f : D \rightarrow \mathbb{R}$ satisfies the partial differential equation

$$\operatorname{div} \left(\frac{\nabla f(z) + 2z^\perp}{|\nabla f(z) + 2z^\perp|} \right) = \frac{3\mathcal{P}(C)}{4\mathcal{V}(C)} \quad (1.10)$$

in $\operatorname{int}(D) \setminus \Sigma(f)$. Here $z = (x, y)$ and $z^\perp = (-y, x)$. The curvature operator in (1.10) has been studied by several authors (besides the previous references, see also e.g. [Pa1], [Pa2], [CHY]). The number

$$H = \frac{3\mathcal{P}(C)}{4\mathcal{V}(C)} \quad (1.11)$$

is the (horizontal) curvature of ∂C , and the equation states that the boundary of a convex isoperimetric set has constant curvature in a weak sense.

If we write $u(z) = \nabla f(z) + 2z^\perp$, then we can interpret equation (1.10) as an equation for the distributional derivative of $u/|u|$, which is a measure. In Section 5, we prove that this measure is absolutely continuous. In other words, we get an improved regularity along horizontal directions for the boundary of convex isoperimetric sets. This result is a corollary of the following regularity theorem for BV vector fields.

Theorem 1.2 (Improved regularity). *Let $\Omega \subseteq \mathbb{R}^2$ be a bounded open set and let $u \in BV(\Omega; \mathbb{R}^2)$ be a vector field. Suppose that:*

- i) There exists $\delta > 0$ such that $|u(z)| \geq \delta$ for $\mathcal{L}^2$ -a.e. $z \in \Omega$;*
- ii) $\operatorname{div} u^\perp \in L^1(\Omega)$;*
- iii) $\operatorname{div} \left(\frac{u}{|u|} \right) \in L^1(\Omega)$.*

Then $u/|u| \in W^{1,1}(\Omega; \mathbb{R}^2)$.

In the reconstruction argument for isoperimetric sets in Section 10, we also need to work with graphs of the form $x = h(y, t)$, where $h : E \rightarrow \mathbb{R}$ is a convex function defined on some bounded convex set $E \subseteq \mathbb{R}^2$ with nonempty interior. For these graphs, the partial differential operator related to the horizontal curvature is a bit more complicated than the one appearing in (1.10) (see equation (4.8)). In Theorem 5.1, we prove a more general version of Theorem 1.2, which includes both curvature operators above as special cases.

The improved regularity of the boundary is the starting point for the geometric characterization of convex isoperimetric sets. A first interesting consequence of Theorem 1.2 is the stability of the curvature with respect to smooth approximations, which is proved in Section 6.

Theorem 1.3 (Stability of the curvature). *Let $\Omega \subseteq \mathbb{R}^2$ be a bounded open set, let $f \in \text{Lip}(\Omega)$ be a Lipschitz function such that $\nabla f \in BV(\Omega; \mathbb{R}^2)$ and denote by $f_\varepsilon \in C^\infty(\Omega_\varepsilon)$ the standard mollification of f , with $\Omega_\varepsilon = \{z \in \Omega \mid \text{dist}(z, \partial\Omega) > \varepsilon\}$. We consider the vector fields*

$$u(z) = \nabla f(z) + 2z^\perp \quad \text{and} \quad u_\varepsilon(z) = \nabla f_\varepsilon(z) + 2z^\perp. \quad (1.12)$$

Assume that:

- i) There exist $\delta > 0$ and $\varepsilon_0 > 0$ such that $|u(z)| \geq \delta$ for $\mathcal{L}^2$ -a.e. $z \in \Omega$ and $|u_\varepsilon(z)| \geq \delta$ for all $z \in \Omega_\varepsilon$, $0 < \varepsilon < \varepsilon_0$;*
- ii) $\text{div}\left(\frac{u}{|u|}\right) \in L^1(\Omega)$.*

Then

$$\lim_{\varepsilon \rightarrow 0} \int_K \left| \text{div}\left(\frac{u_\varepsilon}{|u_\varepsilon|}\right) - \text{div}\left(\frac{u}{|u|}\right) \right| dz = 0 \quad (1.13)$$

for any compact set $K \subseteq \Omega$.

The vector field $v(z) = -u^\perp(z) = 2z - \nabla f^\perp(z)$ belongs to $BV_{\text{loc}}(\text{int}(D); \mathbb{R}^2)$ if $f : D \rightarrow \mathbb{R}$ is convex. Moreover, its distributional divergence is in L^∞ , in fact

$$\text{div } v = 4 \quad \text{in } \text{int}(D). \quad (1.14)$$

Therefore, we can use the theory on the Cauchy Problem for vector fields of bounded variation recently developed by Ambrosio in [A]. This theory extends the work of DiPerna and Lions on the flow of Sobolev vector fields ([DL]) to the BV setting. The bound on the divergence ensures the existence of a unique regular Lagrangian flow $\Phi : K \times [-\varrho, \varrho] \rightarrow D$ ($\varrho > 0$ small enough) relative to v starting from a compact set $K \subseteq \text{int}(D)$. For $\mathcal{L}^2$ -a.e. $z \in K$, the curve $s \mapsto \Phi(z, s)$ is an integral curve of v passing through z at time $s = 0$.

In Sections 7, 8 and 9, we show that $\mathcal{L}^2$ -a.e. integral curve of v is an arc of circle. Since the horizontal lift of the flow Φ foliates the graph of the function f , it follows that the boundary of the convex isoperimetric set is foliated by geodesics.

Theorem 1.4 (Foliation by circles). *Let $C \subseteq \mathbb{H}$ be a convex isoperimetric set with curvature $H > 0$. Then, for $\mathcal{L}^2$ -a.e. $z \in K$, the integral curve $s \mapsto \Phi(z, s)$ is an arc of circle with radius $1/H$.*

The precise version of this result is Theorem 7.2 in Section 7. In Theorem 7.3 we prove an analogous result for graphs of the form $x = h(y, t)$. We give two different proofs of these theorems. The proof in Section 8 is based on the stability property of regular Lagrangian flows and on the stability property (1.13) of the curvature. Here, we describe the proof contained in Section 9, which is based on the Sobolev regularity for $u/|u|$ of Theorem 1.2 and on the regularity of the Lagrangian flow.

The vector field v has a regular Lagrangian flow $\Phi : K \times [-\varrho, \varrho] \rightarrow D$ but is only in $BV_{\text{loc}}(\text{int}(D); \mathbb{R}^2)$. On the other hand, $v/|v|$ is in $W_{\text{loc}}^{1,1}(\text{int}(D); \mathbb{R}^2)$, but its divergence

is only in $L^1_{\text{loc}}(\text{int}(D))$. In order to compute the second order derivative of a generic integral curve of v , we introduce suitable reparameterizations $\gamma(s) = \Phi(z, \tau(s))$. In Theorem 9.1, we show that for any vector field $w \in W^{1,1}(\Omega; \mathbb{R}^2)$ defined in some open neighbourhood Ω of K , the curve $\kappa(s) = w(\gamma(s))$ is in $W^{1,1}$ (for $\mathcal{L}^2$ -a.e. $z \in K$) and moreover

$$\dot{\kappa} = (\nabla w \circ \gamma) \dot{\gamma} \quad \text{in the weak sense.} \quad (1.15)$$

Using the chain rule (1.15) with $w = v/|v|$, we can give a pointwise meaning to equation (1.10) along the flow, and the integral curves of v turn out to be arcs of circles. The proof of Theorem 7.3 given in Section 9 is based on the same technique.

We combine Theorem 7.2 and Theorem 7.3 in Section 10, where we finish the proof of Theorem 1.1.

2 Preliminaries

We can identify the horizontal plane spanned by the vector fields X and Y at the point $p = 0$ with the xy -plane

$$H_0 = \{(z, 0) \in \mathbb{H} \mid z \in \mathbb{C}\}.$$

The yt -plane is the subgroup $\{(z, t) \in \mathbb{H} \mid \text{Re } z = 0\}$. In general, if $p = (z, t) \in \mathbb{H}$, we define the horizontal plane at p as the left translation of H_0 with p , that is

$$H_p = pH_0 = \{(z + z', t + 2\text{Im}z\bar{z}') \in \mathbb{H} \mid z' \in \mathbb{C}\}.$$

The base point p is uniquely determined by H_p . The plane H_p is the boundary of the two halfspaces

$$\begin{aligned} H_p^- &= \{(z', t') \in \mathbb{H} \mid t' \leq t + 2\text{Im}z\bar{z}'\}, \\ H_p^+ &= \{(z', t') \in \mathbb{H} \mid t' \geq t + 2\text{Im}z\bar{z}'\}. \end{aligned} \quad (2.1)$$

We equip the tangent bundle $T\mathbb{H}$ with the left invariant inner product $\langle \cdot, \cdot \rangle$ which makes X , Y and T orthonormal. An absolutely continuous path $\gamma : [0, 1] \rightarrow \mathbb{H}$ is said to be horizontal if $\dot{\gamma}(s) \in \text{span}_{\mathbb{R}}\{X(\gamma(s)), Y(\gamma(s))\}$ for a.e. $s \in [0, 1]$. The curve $\gamma = (\gamma_1, \gamma_2, \gamma_3)$ is horizontal if and only if

$$\gamma_3(s) = \gamma_3(0) + 2 \int_0^s (\dot{\gamma}_1 \gamma_2 - \dot{\gamma}_2 \gamma_1) d\sigma. \quad (2.2)$$

We call the plane curve $\kappa = (\gamma_1, \gamma_2)$ the horizontal projection of γ . If $\kappa = (\gamma_1, \gamma_2)$ is a given plane curve, then a curve $\gamma = (\kappa, \gamma_3)$ with γ_3 given by (2.2) for some $\gamma_3(0)$ is called a horizontal lift of κ .

The sub-Riemannian length of γ is

$$L(\gamma) = \int_0^1 \langle \dot{\gamma}(s), \dot{\gamma}(s) \rangle^{1/2} ds = \int_0^1 |\dot{\kappa}(s)| ds, \quad (2.3)$$

where κ is the horizontal projection of γ and $|\kappa|$ is the Euclidean length of $\kappa \in \mathbb{R}^2$. A path $\gamma : [0, 1] \rightarrow \mathbb{H}$ is said to be admissible for a given pair $(p_0, p_1) \in \mathbb{H} \times \mathbb{H}$ if $\gamma(0) = p_0$, $\gamma(1) = p_1$, and γ is an absolutely continuous, horizontal curve. The Carnot–Carathéodory distance of points $p_0, p_1 \in \mathbb{H}$ is

$$d(p_0, p_1) = \inf \{L(\gamma) \mid \gamma \text{ is admissible for } (p_0, p_1)\}.$$

The open Carnot–Carathéodory ball of radius r centered at $p \in \mathbb{H}$ is denoted by $B_d(p, r)$. The distance d induces the Euclidean topology on $\mathbb{H}$. The metric space $(\mathbb{H}, d)$ is complete, locally compact and geodesic.

Geodesic curves (i.e. length minimizing curves between points) can be computed explicitly, by minimizing the length functional on the right hand side of (2.3) subject to the constraint (2.2) with $s = 1$. Take points $p_0 = 0$ and $p_1 = (z, t) \in \mathbb{H}$. We have the following cases:

- 1) If $t = 0$, the geodesic $\gamma : [0, 1] \rightarrow \mathbb{H}$ connecting 0 to $p_1 = (z, 0)$ is the line segment $\gamma(s) = (sz, 0)$, $s \in [0, 1]$.
- 2) If $t > 0$ and $z \neq 0$, the geodesic connecting 0 to $p_1 = (z, t)$ is the horizontal lift (starting at 0) of the arc of circle from 0 to z in the xy -plane (oriented clockwise), such that the plane region bounded by the arc and by the segment joining 0 to z has area equal to $t/4$. This geodesic is unique.
- 3) If $t > 0$ and $z = 0$, the geodesic from 0 to $p_1 = (0, t)$ is not unique. Take any full circle (oriented clockwise) passing through 0 and with area equal to $t/4$. The horizontal lift of the circle (starting from 0) is the desired geodesic.

The case $t < 0$ is similar and the case $p_0 \neq 0$ is obtained by left translation. If the arc of circle in 2) and 3) has radius $0 < R < +\infty$, we say that the geodesic has curvature $H = 1/R$.

The union of all geodesics joining $(0, -\pi/2)$ to $(0, \pi/2)$ of case 3) bounds the isoperimetric set conjectured by Pansu. The horizontal lift of the plane circle $\kappa(s) = \frac{1}{2}(1 + \cos s, -\sin s)$, $s \in [-\pi, \pi]$, passing through the point $(1, 0, 0) \in \mathbb{H}$ at time $s = 0$ is the curve $\gamma : [-\pi, \pi] \rightarrow \mathbb{H}$

$$\gamma(s) = \frac{1}{2}(1 + \cos s, -\sin s, s + \sin s). \quad (2.4)$$

The third coordinate can be computed using formula (2.2). The curve γ is a geodesic with curvature $H = 2$, starting from $\gamma(-\pi) = (0, 0, -\pi/2)$ and reaching $\gamma(\pi) = (0, 0, \pi/2)$. If $(z, t) = \gamma(s) \in \gamma([-\pi, \pi])$ is a point on the curve, then we have

$$|z| = \left(\frac{1 + \cos s}{2}\right)^{1/2} \quad \text{and} \quad t = \frac{1}{2}(s + \sin s),$$

and we obtain the relation $|t| = \arccos |z| + |z|\sqrt{1 - |z|^2}$. We call the set

$$C_2 = \left\{ (z, t) \in \mathbb{H} \mid |t| < \arccos |z| + |z|\sqrt{1 - |z|^2}, |z| < 1 \right\} \quad (2.5)$$

isoperimetric bubble with curvature $H = 2$. The boundary of C_2 is a compact surface which is obtained by rotation of the generating curve (2.4) around the t -axis. This surface is globally of class C^2 , but fails to be of class C^3 at the characteristic points $(0, \pm\pi/2)$.

We conclude this preliminary section with the following representation formula for perimeter. If $A \subseteq \mathbb{H}$ is a bounded open set such that ∂A is a Lipschitz surface, then

$$\mathcal{P}(A) = \int_{\partial A} \sqrt{(X \cdot \nu)^2 + (Y \cdot \nu)^2} d\mathcal{H}^2, \quad (2.6)$$

where ν is a unit normal to ∂A . Here and in the following, $\cdot$ denotes the standard inner product in $\mathbb{R}^3$ or $\mathbb{R}^2$ (we think of X and Y as vectors in $\mathbb{R}^3$). $\mathcal{H}^2$ is the 2-dimensional Hausdorff measure in $\mathbb{R}^3$ (with respect to the usual Euclidean distance). For a proof of formula (2.6), see [FSSC].

3 Characteristic set of convex sets and convex functions

A set $C \subseteq \mathbb{H}$ is convex if it is convex with respect to the standard convexity of $\mathbb{R}^3$ as a vector space. Note that this notion is invariant with respect to affine transformations of $\mathbb{R}^3$. In particular, it is invariant with respect to left translations, dilations and isometries of $(\mathbb{H}, d)$. Given $p \in \partial C$ and a plane $\pi \subseteq \mathbb{H}$ passing through p , we say that π is a supporting plane for C at p if $\pi \cap \overline{C} \subseteq \partial C$. The property of being a supporting plane for some convex set C is a local property. The characteristic set of C is

$$\Sigma(C) = \{p \in \partial C \mid H_p \text{ is a supporting plane for } C \text{ at } p\},$$

where H_p is the horizontal plane with base point p .

Lemma 3.1. *Let $C \subseteq \mathbb{H}$ be a convex set such that $B_d(p, r_1) \subseteq C \subseteq B_d(p, r_2)$ for some $p \in \mathbb{H}$ and suitable $0 < r_1 < r_2$. Then there exists a sequence $\{C_k\}_{k \in \mathbb{N}}$ of strictly convex sets of class C^∞ such that $B_d(p, r_1/2) \subseteq C_k \subseteq B_d(p, 2r_2)$ for all sufficiently large $k \in \mathbb{N}$ and $C_k \rightarrow C$ as $k \rightarrow \infty$ with respect to the Hausdorff distance.*

Proof. After a left translation, we can assume that $p = 0$. Consider the functions $f_k = \chi_{1/k} * (g + |\cdot|^2/k)$, where χ is a smoothing kernel and g is the homogeneous convex gauge function associated to C . Then each f_k is a smooth, strictly convex function, and the sets $C_k = \{p \in \mathbb{R}^3 \mid f_k(p) \leq 1\}$ have the required properties. $\square$

Lemma 3.2. *If $C \subseteq \mathbb{H}$ is a convex, bounded open set with C^∞ boundary, then $\Sigma(C) \neq \emptyset$.*

Proof. For $p \in \partial C$, let us denote by $V(p) \in T_p \partial C$ the orthogonal projection of the vector field T onto $T_p \partial C$ with respect to the left invariant inner product $\langle \cdot, \cdot \rangle$. Since ∂C is diffeomorphic with the sphere S^2 , there exists $p \in \partial C$ such that $V(p) = 0$. Hence $T_p \partial C = H_p$, and therefore $p \in \Sigma(C)$. $\square$

Lemma 3.3. *If $C \subseteq \mathbb{H}$ is a strictly convex, bounded set, then $\Sigma(C)$ contains at most two points.*

Proof. Let $p_0, p_1 \in \Sigma(C)$ be two characteristic points such that $p_0 \neq p_1$. After a left translation, we can assume that $p_0 = 0$ and $p_1 = (z_1, t_1)$ with $t_1 > 0$. The convex set C is contained between the horizontal planes H_{p_0} and H_{p_1} . By strict convexity of C , the relative interior of the line segment $S = \{(sz_1, st_1) \in \mathbb{H} \mid 0 \leq s \leq 1\}$ connecting p_0 with p_1 is contained in the interior of C . Moreover, letting $p_s = (sz_1, st_1)$, we have

$$\partial C = \{p_0, p_1\} \cup \bigcup_{0 < s < 1} \partial C \cap H_{p_s}.$$

Note that, by strict convexity, $\partial C \cap H_{p_0} = \{p_0\}$ and $\partial C \cap H_{p_1} = \{p_1\}$.

Using the property $p \in H_{p'}$ if and only if $p' \in H_p$, we deduce that if $p \in \partial C \cap H_{p_s}$ with $0 < s < 1$, then $p_s \in H_p$ and H_p is not a supporting plane of C , because p_s belongs to the interior of C . It follows that $\Sigma(C) = \{p_0, p_1\}$. $\square$

Proposition 3.4. *For all $0 < r_1 < r_2$, there is $\varepsilon > 0$, such that, for any convex set $C \subseteq \mathbb{H}$ with $B_d(p, r_1) \subseteq C \subseteq B_d(p, r_2)$ for some $p \in \mathbb{H}$, there exist points $p_0, p_1 \in \Sigma(C)$ with $d(p_0, p_1) \geq \varepsilon$.*

Proof. We first assume that C is a strictly convex set with C^∞ boundary. By Lemmas 3.2 and 3.3, $\Sigma(C)$ contains at least one and at most two points. After a left translation and an isometry, we have $p_0 = 0 \in \Sigma(C)$ and $C \subseteq H_0^+$.

There exist $r > 0$ and a smooth, strictly convex function

$$f : \{z \in \mathbb{R}^2 \mid |z| < r\} \rightarrow \mathbb{R},$$

such that $f(0) = 0$, $f(z) > 0$ for $0 < |z| < r$, and $(z, f(z)) \in \partial C$ for all z .

Denote by V the orthogonal projection of the vector field T onto $T\partial C$ with respect to the left invariant inner product $\langle \cdot, \cdot \rangle$, and let W be the projection of V onto the xy -plane. Here and in the following, when we speak of a projection onto the xy - or yt -plane, it is understood that the projection is orthogonal in the Euclidean sense. Thinking of W as a mapping $W : \{z \in \mathbb{R}^2 \mid |z| < r\} \rightarrow \mathbb{R}^2$, we have

$$W(z) = \frac{\nabla f(z) + 2z^\perp}{1 + |\nabla f(z) + 2z^\perp|^2} \quad \text{for all } z.$$

Observe that $(z, f(z)) \in \Sigma(C)$ if and only if $\nabla f(z) + 2z^\perp = 0$. Hence 0 is the unique zero of W in $\{z \in \mathbb{R}^2 \mid |z| < r\}$, provided that $r > 0$ is small enough. Let

$$K := \{z \in \mathbb{R}^2 \mid |z| \leq r/2\}.$$

Let $\partial_r f$ denote the radial derivative of f . By strict convexity of f , the mapping $F : \partial K \rightarrow S^1$ given by

$$F(z) = \frac{\nabla f(z) + 2z^\perp}{|\nabla f(z) + 2z^\perp|}$$

satisfies

$$F(z) \cdot \frac{z}{|z|} = \frac{\partial_r f(z)}{|\nabla f(z) + 2z^\perp|} > 0 \quad \text{on } \partial K.$$

Hence F and the Gauss map $G : \partial K \rightarrow S^1$ are homotopic. It follows that

$$\text{index}(V, 0) = \text{index}(W, 0) = \text{degree}(F, \partial K) = \text{degree}(G, \partial K) = 1. \quad (3.1)$$

If V had only one zero on ∂C , then the Poincaré–Hopf index theorem would give $\text{index}(V, 0) = \chi(C) = \chi(S^2) = 2$, where χ denotes the Euler–Poincaré characteristic. In view of (3.1), this is not possible.

Thus $\Sigma(C) = \{p_0, p_1\}$ consists of exactly two points. We can assume that $p_0 = 0$. We show that $d(p_0, p_1) \geq \varepsilon$ for some $\varepsilon > 0$. There exists an interior point $p = (z, t) \in C$ such that $|z| \leq \alpha$ and $t \geq \beta$ for some $\alpha = \alpha(r_2) < +\infty$ and $\beta = \beta(r_1, r_2) > 0$. Let $S = \{(sz, st) \mid 0 \leq s \leq 1\}$ be the line segment connecting $p_0 = 0 \in \Sigma(C)$ with p . As in the proof of Lemma 3.3, we have $\Sigma(C) \cap H_{(sz, st)} = \emptyset$ for all $0 < s \leq 1$. Therefore

$$\Sigma(C) \cap H_p^- = \{0\}.$$

Moreover, $B_d(0, \varepsilon) \subseteq H_p^-$ for some $\varepsilon > 0$ depending on α and β .

Now assume that C is only convex. By Lemma 3.1, there exists a sequence $\{C_k\}_{k \in \mathbb{N}}$ of strictly convex sets of class C^∞ such that $B_d(p, r_1/2) \subseteq C_k \subseteq B_d(p, 2r_2)$ for all $k \in \mathbb{N}$ and $C_k \rightarrow C$ with respect to the Hausdorff distance. From the first part of the proof, we know that there exist sequences $\{p_{k,0}\}_{k \in \mathbb{N}}$ and $\{p_{k,1}\}_{k \in \mathbb{N}}$ with $p_{k,0}, p_{k,1} \in \Sigma(C_k)$ and $d(p_{k,0}, p_{k,1}) \geq \varepsilon > 0$ for all $k \in \mathbb{N}$. Passing to subsequences and relabelling if necessary, we can assume that these sequences converge to limit points p_0, p_1 with the desired properties. $\square$

Definition 3.1. A line ℓ is horizontal if $\ell \subseteq H_p$ for one (equivalently: all) $p \in \ell$. A horizontal segment is a segment of a horizontal line.

Theorem 3.5 (Characteristic set). *If $C \subseteq \mathbb{H}$ is a convex open set, then $\Sigma(C)$ admits the decomposition $\Sigma(C) = \Sigma^- \cup \Sigma^+$, where Σ^-, Σ^+ are two closed, disjoint, horizontal segments (possibly points). Moreover, if $0 \in \text{int}(C)$, then H_0 separates Σ^- from Σ^+ .*

Proof. Without loss of generality, we can assume $0 \in \text{int}(C)$. We define the convex sets

$$C^- = C \cap H_0^- \quad \text{and} \quad C^+ = C \cap H_0^+.$$

If $p \in \partial C^- \cap H_0 = \partial C^+ \cap H_0$ with $p \neq 0$, then $0 \in H_p \neq H_0$ and consequently $H_p \cap \text{int}(C^\pm) \neq \emptyset$. It follows that $\Sigma(C^\pm) \cap H_0 = \{0\}$, and, by Corollary 3.4, there exist points p^-, p^+ such that $p^\pm \in \Sigma(C^\pm) \cap \text{int}(H_0^\pm)$. The relative interior of the line segment connecting 0 with p^- , respectively p^+ , is contained in the interior of C^- , respectively C^+ . Then the same reasoning as in the proof of Lemma 3.3 gives

$$\Sigma(C^-) = \{0\} \cup (\Sigma(C^-) \cap H_{p^-}) \quad \text{and} \quad \Sigma(C^+) = \{0\} \cup (\Sigma(C^+) \cap H_{p^+}). \quad (3.2)$$

Let $p \in \Sigma(C^-) \cap H_{p^-}$ with $p \neq p^-$. Since (3.2) must hold with p instead of p^- , it follows that $\Sigma(C^-) \cap H_{p^-}$ is contained in a horizontal segment. By compactness

and convexity, there exists a closed, bounded, horizontal segment Σ^- such that $\Sigma(C^-) \cap H_{p^-} = \Sigma^-$. Similarly, we can show that $\Sigma(C^+) \cap H_{p^+} = \Sigma^+$ for some closed, bounded, horizontal segment Σ^+ . Neither Σ^- nor Σ^+ intersects H_0 .

Finally, we show that $\Sigma(C) = \Sigma^- \cup \Sigma^+$. $\Sigma(C) \cap H_0 = \emptyset$ implies $\Sigma(C) \subseteq \Sigma^- \cup \Sigma^+$. On the other hand, $p \in \Sigma^- \cup \Sigma^+$ implies $p \in \Sigma(C)$ (locality). $\square$

We recall some definitions concerning convex functions. Let $\Omega \subseteq \mathbb{R}^2$ be a convex open set and let $f : \Omega \rightarrow \mathbb{R}$ be a convex function. The subdifferential $\partial f(q)$ of f at $q \in \Omega$ is the (nonempty) set

$$\partial f(q) = \{q^* \in \mathbb{R}^2 \mid f(q') - f(q) \geq q^* \cdot (q' - q) \text{ for all } q' \in \Omega\}. \quad (3.3)$$

The property of belonging to the subdifferential is a local property. It is well known that f is locally Lipschitz continuous in Ω , that $\partial f(q) = \{\nabla f(q)\}$ if and only if f is differentiable at $q \in \Omega$ with gradient $\nabla f(q)$ and that $\nabla f \in BV_{\text{loc}}(\Omega; \mathbb{R}^2)$. Moreover, we have the following

Lemma 3.6. *Let $\Omega \subseteq \mathbb{R}^2$ be a convex open set, let $f : \Omega \rightarrow \mathbb{R}$ be a convex function and let $\{f_k\}_{k \in \mathbb{N}}$ be a sequence of convex functions $f_k : \Omega \rightarrow \mathbb{R}$ which converges to f locally uniformly in Ω . Given a compact set $K \subseteq \Omega$, sequences $\{q_k\}_{k \in \mathbb{N}}$ in K and $\{q_k^*\}_{k \in \mathbb{N}}$ with $q_k^* \in \partial f_k(q_k)$ for all $k \in \mathbb{N}$, there exist $q \in K$, $q^* \in \partial f(q)$ and subsequences $\{q_{k_l}\}_{l \in \mathbb{N}}$, $\{q_{k_l}^*\}_{l \in \mathbb{N}}$, such that $q_{k_l} \rightarrow q$ and $q_{k_l}^* \rightarrow q^*$.*

Let C be a bounded, convex compact set in $\mathbb{H}$ with nonempty interior. The projection D of C onto the xy -plane

$$D = \{z \in \mathbb{R}^2 \mid \text{there is } t \in \mathbb{R} \text{ such that } (z, t) \in C\}$$

is a convex compact set in $\mathbb{R}^2$, and there are functions $f, g : D \rightarrow \mathbb{R}$, with f convex and g concave, such that $C = \{(z, t) \in \mathbb{H} \mid z \in D, f(z) \leq t \leq g(z)\}$. We have $f < g$ in $\text{int}(D)$. If $z \in \partial D$, then $f(z) \leq g(z)$, possibly $f(z) < g(z)$.

Definition 3.2. The characteristic set $\Sigma(f)$ of f is the set of points $z \in D$ such that the horizontal plane H_p at the point $p = (z, f(z)) \in \mathbb{R}^2 \times \mathbb{R}$ is a supporting plane for C .

Remark 3.1. A computation shows that

$$z \in \Sigma(f) \cap \text{int}(D) \iff -2z^\perp \in \partial f(z). \quad (3.4)$$

Proposition 3.7 (Lower bounds I). *Let $\Omega \subseteq \text{int}(D)$ be an open set and let $\{f_\varepsilon\}_{\varepsilon > 0}$ be a family of smooth convex functions in Ω such that $f_\varepsilon \rightarrow f$ locally uniformly and $\nabla f_\varepsilon \rightarrow \nabla f$ in $L^1_{\text{loc}}(\Omega; \mathbb{R}^2)$ as $\varepsilon \downarrow 0$. Then, for any compact set $K \subseteq \Omega \setminus \Sigma(f)$, there are constants $\delta > 0$ and $\varepsilon_0 > 0$ such that*

$$|\nabla f_\varepsilon(z) + 2z^\perp| \geq \delta \quad (3.5)$$

for all $0 < \varepsilon < \varepsilon_0$ and all $z \in K$. Consequently,

$$|\nabla f(z) + 2z^\perp| \geq \delta \quad (3.6)$$

for $\mathcal{L}^2$ -a.e. $z \in K$.

Proof. Suppose by contradiction that the pair (δ, ε_0) does not exist. Then, by Lemma 3.6, there are sequences $\{f_{\varepsilon_k}\}_{k \in \mathbb{N}}$, $\{z_k\}_{k \in \mathbb{N}}$ and $\{\nabla f_{\varepsilon_k}(z_k)\}_{k \in \mathbb{N}}$ such that $z_k \rightarrow z \in K$ and $\nabla f_{\varepsilon_k}(z_k) \rightarrow -2z^\perp \in \partial f(z)$. This implies $z \in \Sigma(f)$ by (3.4), contradicting the assumption $K \cap \Sigma(f) = \emptyset$. $\square$

Similarly, the projection E of C onto the yt -plane

$$E = \{\zeta \in \mathbb{R}^2 \mid \text{there is } x \in \mathbb{R} \text{ such that } (x, \zeta) \in C\}$$

is a convex compact set in $\mathbb{R}^2$ and there are functions $h, k : E \rightarrow \mathbb{R}$, with h convex and k concave, such that $C = \{(x, \zeta) \in \mathbb{H} \mid \zeta \in E, h(\zeta) \leq x \leq k(\zeta)\}$. We have $h < k$ in $\text{int}(E)$. If $\zeta \in \partial E$, then $h(\zeta) \leq k(\zeta)$, possibly $h(\zeta) < k(\zeta)$.

Definition 3.3. The characteristic set $\Sigma(h)$ of h is the set of points $\zeta \in E$ such that the horizontal plane H_p at the point $p = (h(\zeta), \zeta) \in \mathbb{R} \times \mathbb{R}^2$ is a supporting plane for C .

Remark 3.2. A computation shows that

$$\zeta = (y, t) \in \Sigma(h) \cap \text{int}(E) \iff y \neq 0 \text{ and } (h(\zeta)/y, 1/(2y)) \in \partial h(\zeta). \quad (3.7)$$

Proposition 3.8 (Lower bounds II). *Let $\Omega \subseteq \text{int}(E)$ be an open set and let $\{h_\varepsilon\}_{\varepsilon > 0}$ be a family of smooth convex functions in Ω such that $h_\varepsilon \rightarrow h$ locally uniformly and $\nabla h_\varepsilon \rightarrow \nabla h$ in $L^1_{\text{loc}}(\Omega; \mathbb{R}^2)$ as $\varepsilon \downarrow 0$. Then, for any compact set $K \subseteq \Omega \setminus \Sigma(h)$, there are constants $\delta > 0$ and $\varepsilon_0 > 0$, such that*

$$|(\partial_y h_\varepsilon - 2h_\varepsilon \partial_t h_\varepsilon, 1 - 2y \partial_t h_\varepsilon)| \geq \delta \quad (3.8)$$

in K for all $0 < \varepsilon < \varepsilon_0$. Consequently,

$$|(\partial_y h - 2h \partial_t h, 1 - 2y \partial_t h)| \geq \delta \quad (3.9)$$

$\mathcal{L}^2$ -a.e. in K .

Proof. For some $\varepsilon_0 > 0$ the functions h_ε , $0 < \varepsilon < \varepsilon_0$, are uniformly bounded and uniformly Lipschitz continuous in some open neighbourhood of K . Therefore, if $r > 0$ is sufficiently small, we have

$$|1 - 2y \partial_t h_\varepsilon| \geq 1/2$$

in $K \cap \{|y| \leq r\}$ for $0 < \varepsilon < \varepsilon_0$. By (3.7) and Lemma 3.6, there exists $\delta_1 > 0$ such that

$$|(\partial_y h_\varepsilon - h_\varepsilon/y, \partial_t h_\varepsilon - 1/(2y))| \geq \delta_1$$

in $K \cap \{|y| \geq r\}$ for $0 < \varepsilon < \varepsilon_0$, with a possibly smaller ε_0 . Hence there exists $\delta_2 > 0$ such that the condition $|\partial_t h_\varepsilon - 1/(2y)| \leq \delta_2$ implies $|\partial_y h_\varepsilon - h_\varepsilon/y| \geq \delta_1/2$ and $|2h_\varepsilon(\partial_t h_\varepsilon - 1/(2y))| \leq \delta_1/4$, whence

$$|\partial_y h_\varepsilon - 2h_\varepsilon \partial_t h_\varepsilon| \geq |\partial_y h_\varepsilon - h_\varepsilon/y| - |2h_\varepsilon(\partial_t h_\varepsilon - 1/(2y))| \geq \delta_1/4.$$

The claim follows with $\delta = \min\{1/2, \delta_1/4, 2r\delta_2\}$. $\square$

We conclude this section with the following observation.

Lemma 3.9 (Uniqueness). *Let $C \subseteq \mathbb{H}$ be a convex compact set with nonempty interior. Suppose that there exist two geodesics contained in ∂C and with curvature $H > 0$ passing through a point $p \in \partial C$. Then they coincide in a neighbourhood of p .*

Proof. We can assume $p = 0$. Let $L > 0$ and let $\gamma_0, \gamma_1 : [-L, L] \rightarrow \partial C$ be geodesics parameterized by arc length, with curvature $H > 0$ and such that $\gamma_0(0) = \gamma_1(0) = 0$. If $\dot{\gamma}_0(0) = \dot{\gamma}_1(0)$, then $\gamma_0 = \gamma_1$. Assume by contradiction that $\dot{\gamma}_0(0) \neq \dot{\gamma}_1(0)$. Then H_0 is a supporting plane for C . (This is true also in the case $\dot{\gamma}_0(0) = -\dot{\gamma}_1(0)$). But this is not possible, because $\gamma_0(s) \in \{t > 0\}$ for $s \in (0, L]$ and $\gamma_0(s) \in \{t < 0\}$ for $s \in [-L, 0)$. $\square$

4 Curvature equations for convex isoperimetric sets

We derive partial differential equations for certain vector fields built from functions which parameterize the boundary of convex isoperimetric sets. These equations state that the boundary of a convex isoperimetric set $C \subseteq \mathbb{H}$ has constant horizontal curvature. We study two different curvature equations: the equation for graphs of the form $t = f(x, y)$ and the equation for graphs of the form $x = h(y, t)$. We call the number $H = 3\mathcal{P}(C)/4\mathcal{V}(C)$ the horizontal curvature of C (curvature for short).

Graphs of the form $t = f(x, y)$. In this subsection, we denote the elements of $\mathbb{H} = \mathbb{R}^2 \times \mathbb{R}$ by (z, t) with $t \in \mathbb{R}$ and $z = (x, y) \in \mathbb{R}^2$. We write $z^\perp = (-y, x)$. Let C be a convex set in $\mathbb{H}$ of the form

$$C = \{(z, t) \in \mathbb{H} \mid z \in D, f(z) \leq t \leq g(z)\}, \quad (4.1)$$

where $D \subseteq \mathbb{R}^2$ is a convex compact set in the plane with nonempty interior, and $-g, f : D \rightarrow \mathbb{R}$ are convex functions.

Proposition 4.1 (Curvature equation I). *If $C \subseteq \mathbb{H}$ is a convex isoperimetric set with perimeter $\mathcal{P}(C)$ and volume $\mathcal{V}(C)$, then the function $f : D \rightarrow \mathbb{R}$ satisfies in distributional sense the partial differential equation*

$$\operatorname{div} \left(\frac{\nabla f(z) + 2z^\perp}{|\nabla f(z) + 2z^\perp|} \right) = H \quad (4.2)$$

in $\operatorname{int}(D) \setminus \Sigma(f)$, with $H = 3\mathcal{P}(C)/4\mathcal{V}(C)$.

Proof. By Theorem 3.5, the characteristic set $\Sigma(f) \subseteq D$ is a union of two (possibly empty) closed line segments. Let $\varphi \in C_c^\infty(\operatorname{int}(D) \setminus \Sigma(f))$ and for $\varepsilon \in \mathbb{R}$, consider the set

$$C_\varepsilon = \{(z, t) \in \mathbb{H} \mid z \in D, f(z) + \varepsilon\varphi(z) \leq t \leq g(z)\}.$$

The set C_ε is nonempty when $|\varepsilon|$ is sufficiently small. We write $\mathcal{P}(\varepsilon) = \mathcal{P}(C_\varepsilon)$, $\mathcal{V}(\varepsilon) = \mathcal{V}(C_\varepsilon)$ and $\mathcal{I}(\varepsilon) = \mathcal{P}(\varepsilon)^4/\mathcal{V}(\varepsilon)^3$.

If the set C minimizes the isoperimetric ratio $\mathcal{I}(C) = \mathcal{P}(C)^4/\mathcal{V}(C)^3$, then the function $\mathcal{I}(\varepsilon)$ has a minimum at $\varepsilon = 0$, and therefore we have the equation

$$\mathcal{I}'(0) = \frac{\mathcal{P}^3}{\mathcal{V}^4} (4\mathcal{P}'\mathcal{V} - 3\mathcal{P}\mathcal{V}') \Big|_{\varepsilon=0} = 0. \quad (4.3)$$

There exists $\varepsilon_0 > 0$ such that

$$\mathcal{V}'(\varepsilon) = - \int_D \varphi(z) dz \quad \text{for } |\varepsilon| < \varepsilon_0. \quad (4.4)$$

Moreover, denoting by

$$S_\varepsilon = \{(z, f(z) + \varepsilon\varphi(z)) \in \mathbb{H} \mid z \in D\}$$

the graph of the function $f + \varepsilon\varphi$ and by

$$\nu_\varepsilon = \frac{(\nabla f + \varepsilon \nabla \varphi, -1)}{(1 + |\nabla f + \varepsilon \nabla \varphi|^2)^{1/2}}$$

the exterior unit normal to S_ε (defined $\mathcal{H}^2$ -a.e. on S_ε), from the Heisenberg Area Formula (2.6) and from the standard Area Formula for graphs of functions in Euclidean spaces, we find

$$\begin{aligned} \mathcal{P}'(\varepsilon) &= \frac{d}{d\varepsilon} \int_{S_\varepsilon} \sqrt{(X \cdot \nu_\varepsilon)^2 + (Y \cdot \nu_\varepsilon)^2} d\mathcal{H}^2 \\ &= \frac{d}{d\varepsilon} \int_D |\nabla f(z) + \varepsilon \nabla \varphi(z) + 2z^\perp| dz. \end{aligned} \quad (4.5)$$

By Proposition 3.7, for each compact set $K \subseteq \text{int}(D) \setminus \Sigma(f)$, there is $\delta > 0$ such that $|\nabla f(z) + 2z^\perp| \geq \delta$ for $\mathcal{L}^2$ -a.e. $z \in K$. Then $|\nabla f(z) + \varepsilon \nabla \varphi(z) + 2z^\perp|$ is essentially bounded away from 0 in a neighbourhood of $\text{spt}(\varphi)$ when $|\varepsilon|$ is small enough and we can interchange derivative and integral in the second line of (4.5). At $\varepsilon = 0$ we obtain

$$\mathcal{P}'(0) = \int_D \frac{\nabla f(z) + 2z^\perp}{|\nabla f(z) + 2z^\perp|} \cdot \nabla \varphi(z) dz. \quad (4.6)$$

From (4.3), (4.4) and (4.6) we find

$$4\mathcal{V}(C) \int_D \frac{\nabla f(z) + 2z^\perp}{|\nabla f(z) + 2z^\perp|} \cdot \nabla \varphi(z) dz = -3\mathcal{P}(C) \int_D \varphi(z) dz$$

for arbitrary $\varphi \in C_c^\infty(\text{int}(D) \setminus \Sigma(f))$. This is (4.2). $\square$

Graphs of the form $x = h(y, t)$. In this subsection, we denote the elements of $\mathbb{H} = \mathbb{R} \times \mathbb{R}^2$ by (x, ζ) with $x \in \mathbb{R}$ and $\zeta = (y, t) \in \mathbb{R}^2$. Let C be a convex set in $\mathbb{H}$ of the form

$$C = \{(x, \zeta) \in \mathbb{H} \mid \zeta \in E, h(\zeta) \leq x \leq k(\zeta)\}, \quad (4.7)$$

where $E \subseteq \mathbb{R}^2$ is a convex compact set in the plane with nonempty interior and $-k, h : E \rightarrow \mathbb{R}$ are convex functions.

Proposition 4.2 (Curvature equation II). *If $C \subseteq \mathbb{H}$ is a convex isoperimetric set with perimeter $\mathcal{P}(C)$ and volume $\mathcal{V}(C)$, then the function $h : E \rightarrow \mathbb{R}$ satisfies in distributional sense the partial differential equation*

$$(\partial_y - 2h\partial_t)\left(\frac{u_1}{|u|}\right) - 2y\partial_t\left(\frac{u_2}{|u|}\right) = H \quad (4.8)$$

in $\text{int}(E) \setminus \Sigma(h)$, with $H = 3\mathcal{P}(C)/4\mathcal{V}(C)$ and

$$u = (u_1, u_2) = (h_y - 2hh_t, 1 - 2yh_t). \quad (4.9)$$

Proof. By Theorem 3.5, the characteristic set $\Sigma(h)$ is the union of two (possibly empty) closed line segments. We have $u \in BV_{\text{loc}}(\text{int}(E); \mathbb{R}^2)$ and, by Proposition 3.8, for each compact set $K \subseteq (\text{int}(E) \setminus \Sigma(h))$, there exists $\delta > 0$ such that $|u| \geq \delta$ $\mathcal{L}^2$ -a.e. in K . Hence $u/|u| \in BV_{\text{loc}}(\text{int}(E) \setminus \Sigma(h); \mathbb{R}^2)$.

Let $\varphi \in C_c^\infty(\text{int}(E) \setminus \Sigma(h))$. For any $\varepsilon \in \mathbb{R}$ consider the set

$$C_\varepsilon = \{(x, \zeta) \in \mathbb{H} \mid \zeta \in E, h(\zeta) + \varepsilon\varphi(\zeta) \leq x \leq k(\zeta)\},$$

and let $\mathcal{P}(\varepsilon) = \mathcal{P}(C_\varepsilon)$, $\mathcal{V}(\varepsilon) = \mathcal{V}(C_\varepsilon)$ and $\mathcal{I}(\varepsilon) = \mathcal{P}(\varepsilon)^4/\mathcal{V}(\varepsilon)^3$.

Denoting by $S_\varepsilon = \{(h(\zeta) + \varepsilon\varphi(\zeta), \zeta) \in \mathbb{H} \mid \zeta \in E\}$ the graph of the function $h + \varepsilon\varphi$ and by

$$\nu_\varepsilon = \frac{(-1, \nabla h + \varepsilon \nabla \varphi)}{(1 + |\nabla h + \varepsilon \nabla \varphi|^2)^{1/2}}$$

the exterior unit normal to S_ε (defined $\mathcal{H}^2$ -a.e. on S_ε), from the Heisenberg Area Formula (2.6) and from the standard Area Formula, we get

$$\begin{aligned} \mathcal{P}'(\varepsilon) &= \frac{d}{d\varepsilon} \int_{S_\varepsilon} \sqrt{(X \cdot \nu_\varepsilon)^2 + (Y \cdot \nu_\varepsilon)^2} d\mathcal{H}^2 \\ &= \frac{d}{d\varepsilon} \int_E \left((1 - 2y(h_t + \varepsilon\varphi_t))^2 + ((h_y + \varepsilon\varphi_y) - 2(h + \varepsilon\varphi)(h_t + \varepsilon\varphi_t))^2 \right)^{1/2} d\zeta. \end{aligned} \quad (4.10)$$

Let $\delta_1 > 0$ such that $(h_y - 2hh_t)^2 + (1 - 2yh_t)^2 \geq \delta_1^2$ $\mathcal{L}^2$ -a.e. in a neighbourhood $\Omega \subseteq \text{int}(E) \setminus \Sigma(h)$ of $\text{spt}(\varphi)$. We can find $\varepsilon_0 > 0$ such that the integrand in (4.10) is larger than $\delta_1/2$ $\mathcal{L}^2$ -a.e. in Ω whenever $|\varepsilon| < \varepsilon_0$. Then we can differentiate under the integral sign and we obtain

$$\mathcal{P}'(0) = \int_E \frac{(h_y - 2hh_t)(\varphi_y - 2(\varphi h)_t) - 2y\varphi_t(1 - 2yh_t)}{\sqrt{(h_y - 2hh_t)^2 + (1 - 2yh_t)^2}} d\zeta$$

at $\varepsilon = 0$. A formal integration by parts yields

$$\mathcal{P}'(0) = - \int_E \left((\partial_y - 2h\partial_t)\left(\frac{u_1}{|u|}\right) - 2y\partial_t\left(\frac{u_2}{|u|}\right) \right) \varphi d\zeta. \quad (4.11)$$

If the set C minimizes the isoperimetric ratio $\mathcal{I}(C) = \mathcal{P}(C)^4/\mathcal{V}(C)^3$, then, as in (4.3), we have $4\mathcal{P}'(0)\mathcal{V}(0) - 3\mathcal{P}(0)\mathcal{V}'(0) = 0$. From this equation, from (4.11) and from

$$\mathcal{V}'(0) = - \int_E \varphi(\zeta) d\zeta,$$

we get

$$4\mathcal{V}(C) \int_E \left((\partial_y - 2h\partial_t) \left(\frac{u_1}{|u|} \right) - 2y\partial_t \left(\frac{u_2}{|u|} \right) \right) \varphi d\zeta = 3\mathcal{P}(C) \int_E \varphi d\zeta$$

for arbitrary $\varphi \in C_c^\infty(\text{int}(E) \setminus \Sigma(h))$. This is (4.8). $\square$

5 Improved regularity of the boundary

We recall some general properties of BV vector fields. We refer the reader to [AFP], chapter 3, for a detailed discussion and proofs. Let $\Omega \subseteq \mathbb{R}^2$ be an open set. A vector field $u \in BV(\Omega; \mathbb{R}^2)$ has approximate limit $\bar{u}(q) \in \mathbb{R}^2$ at the point $q \in \Omega$ if

$$\lim_{r \downarrow 0} \int_{B(q,r)} |u - \bar{u}(q)| d\mathcal{L}^2 = 0. \quad (5.1)$$

Here and in the following, $B(q, r)$ denotes the open Euclidean ball of radius r centered at q . The approximate limit (if it exists) is uniquely determined by (5.1). The approximate discontinuity set of u is the set S_u of points $q \in \Omega$ at which u has no approximate limit. The set S_u is a Borel set with vanishing $\mathcal{L}^2$ -measure. A point $q \in \Omega$ is an approximate jump point of u if there exist $u^+(q) \in \mathbb{R}^2$, $u^-(q) \in \mathbb{R}^2$ and $\nu_u(q) \in S^1$ such that $u^+(q) \neq u^-(q)$ and

$$\lim_{r \downarrow 0} \int_{B^\pm(q,r)} |u - u^\pm(q)| d\mathcal{L}^2 = 0, \quad (5.2)$$

where

$$B^\pm(q, r) = \{q' \in B(q, r) \mid \pm(q' - q) \cdot \nu_u(q) > 0\}.$$

The triplet $(u^+(q), u^-(q), \nu_u(q))$ is uniquely determined by (5.2) up to multiplication of $\nu_u(q)$ with -1 and permutation of $u^+(q)$ and $u^-(q)$. The set J_u of approximate jump points is an $\mathcal{H}^1$ -rectifiable Borel subset of S_u with $\mathcal{H}^1(S_u \setminus J_u) = 0$. Moreover, there exist Borel functions $u^+, u^- : J_u \rightarrow \mathbb{R}^2$ and $\nu_u : J_u \rightarrow S^1$ such that (5.2) holds at every $q \in J_u$. The precise representative $u^* : \Omega \rightarrow \mathbb{R}^2$ of u is defined as follows:

$$u^*(q) = \begin{cases} \bar{u}(q) & q \in \Omega \setminus S_u, \\ \frac{1}{2}(u^+(q) + u^-(q)) & q \in J_u, \\ 0 & q \in S_u \setminus J_u. \end{cases} \quad (5.3)$$

By the Lebesgue density theorem, $u(q) = u^*(q)$ for $\mathcal{L}^2$ -a.e. $q \in \Omega$. Notice that $|u| \geq \delta > 0$ $\mathcal{L}^2$ -a.e. in Ω implies

$$|\bar{u}(q)| \geq \delta \quad \text{for all } q \in \Omega \setminus S_u \quad \text{and} \quad |u^\pm(q)| \geq \delta \quad \text{for all } q \in J_u. \quad (5.4)$$

By the Riesz representation theorem and the Lebesgue decomposition theorem, there exist vector-valued Radon measures $D^a u$ and $D^s u$ on Ω such that $Du = D^a u + D^s u$ with $D^a u \ll \mathcal{L}^2$ and $D^s u \perp \mathcal{L}^2$. Then we have the decomposition

$$Du = D^a u + D^c u + D^j u,$$

where

$$D^a u = \nabla u \mathcal{L}^2 \quad (5.5)$$

with $\nabla u \in L^1(\Omega; M^{2 \times 2})$ and $M^{2 \times 2}$ is the space of 2×2 matrices with real entries. We have $u \in W^{1,1}(\Omega; \mathbb{R}^2)$ if and only if $D^s u$ vanishes. The Cantor part $D^c u$ and the jump part $D^j u$ of $D^s u$ are respectively

$$D^c u = D^s u \llcorner \Omega \setminus S_u \quad \text{and} \quad D^j u = D^s u \llcorner S_u.$$

By the representation theorem of Federer–Vol’pert and by the rank one theorem of Alberti, the measures $D^c u$ and $D^j u$ admit the representations

$$D^j u = (u^+ - u^-) \otimes \nu_u \mathcal{H}^1 \llcorner J_u \quad \text{and} \quad D^c u = \eta \otimes \xi |D^c u|, \quad (5.6)$$

where $|D^c u|$ is the total variation of $D^c u$ and $\eta, \xi : \Omega \rightarrow S^1$ are suitable Borel maps. The measure $|D^c u|$ is absolutely continuous with respect to $\mathcal{H}^1$.

Finally, we need the chain rule for BV functions. Let $F \in C^1(\mathbb{R}^2; \mathbb{R}^2)$ be a mapping such that $F(0) = 0$ and $|\nabla F| \in L^\infty(\mathbb{R}^2)$. The function $v = F \circ u$ is in $BV(\Omega; \mathbb{R}^2)$ and $v \in L^\infty(\Omega; \mathbb{R}^2)$ if $u \in L^\infty(\Omega; \mathbb{R}^2)$. Moreover, we have

$$D^a v = \nabla F(u^*) \nabla u \mathcal{L}^2 \quad \text{and} \quad D^c v = (\nabla F(u^*) \eta) \otimes \xi |D^c u|. \quad (5.7)$$

A point $q \in \Omega$ belongs to J_v if and only if $q \in J_u$ and $F(u^+(q)) \neq F(u^-(q))$. What’s more

$$D^j v = (F(u^+) - F(u^-)) \otimes \nu_u \mathcal{H}^1 \llcorner J_u. \quad (5.8)$$

We are now in a position to prove our regularity theorem for vector fields with bounded variation arising from the parameterization of the boundary of convex isoperimetric sets. Let $\Omega \subseteq \mathbb{R}^2$ be an open set and let $a, b \in C(\Omega)$ be continuous functions. We consider the differential operator $\mathcal{M}$ defined by

$$\mathcal{M}u = (\partial_1 - a\partial_2)u_1 + b\partial_2 u_2. \quad (5.9)$$

We have $\mathcal{M} = \text{div}$ when $a = 0$ and $b = 1$. On the other hand, when $a = 2h$ and $b = -2y$, $\mathcal{M}$ is the operator appearing in the left hand side of (4.8).

Theorem 5.1 (Improved regularity). *Let $\Omega \subseteq \mathbb{R}^2$ be a bounded open set, let $u = (u_1, u_2) \in BV(\Omega; \mathbb{R}^2)$ be a vector field, and let $a, b \in C(\Omega)$ be continuous functions such that $b \neq 0$ in Ω . Assume that:*

- i) *There exists $\delta > 0$ such that $|u(q)| \geq \delta$ for $\mathcal{L}^2$ -a.e. $q \in \Omega$;*
- ii) *$\mathcal{M}u^\perp \in L^1(\Omega)$, where $u^\perp = (-u_2, u_1)$;*
- iii) *$\mathcal{M}\left(\frac{u}{|u|}\right) \in L^1(\Omega)$.*

Then $u/|u| \in W^{1,1}(\Omega; \mathbb{R}^2)$ and there exists a function $\mu : J_u \rightarrow (0, +\infty)$, such that $u^- = \mu u^+$ on J_u .

Proof. We claim that both the jump and the Cantor part of $D(u/|u|)$ vanish. Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a smooth mapping such that

$$F(0) = 0, \quad F(q) = \frac{q}{|q|} \text{ for } |q| \geq \frac{\delta}{2} \quad \text{and} \quad |\nabla F| \in L^\infty(\mathbb{R}^2). \quad (5.10)$$

Notice that, if $|q| > \delta/2$, the derivative of F has the form

$$\nabla F(q) = \frac{1}{|q|^3} q^\perp \otimes q^\perp. \quad (5.11)$$

By assumption i), the vector field $v = F \circ u$ is in $BV(\Omega; \mathbb{R}^2)$. By (5.4) and (5.8), its jump set is

$$J_v = \{q \in J_u \mid u^+(q)/|u^+(q)| \neq u^-(q)/|u^-(q)|\}$$

and its jump part is

$$D^j v = \left(\frac{u^+}{|u^+|} - \frac{u^-}{|u^-|} \right) \otimes \nu_u \mathcal{H}^1 \llcorner J_u. \quad (5.12)$$

Analogously, if $D^c u = \eta \otimes \xi |D^c u|$ for Borel functions $\eta, \xi : \Omega \rightarrow S^1$, then

$$D^c v = (\nabla F(u^*) \eta) \otimes \xi |D^c u| \quad (5.13)$$

by (5.7).

By assumption ii), the part of the measure

$$-\partial_1 u_2 + b\partial_2 u_1 + a\partial_2 u_2$$

concentrated on J_u vanishes. From the formula for $D^j u$ in (5.6), we can compute $D_k^j u_l$ for $k, l = 1, 2$, and we obtain

$$\left((u_2^- - u_2^+), b(u_1^+ - u_1^-) + a(u_2^+ - u_2^-) \right) \cdot \nu_u = 0 \quad (5.14)$$

$\mathcal{H}^1$ -a.e. on J_u .

By assumption iii), the part of the measure

$$(\partial_1 - a\partial_2)(u_1/|u|) + b\partial_2(u_2/|u|)$$

concentrated on J_v vanishes. Thus, using (5.12), we can compute $D_k^j(u_l/|u|)$ for $k, l = 1, 2$, and we get

$$\left(\frac{u_1^+}{|u^+|} - \frac{u_1^-}{|u^-|}, \frac{au_1^- - bu_2^-}{|u^-|} - \frac{au_1^+ - bu_2^+}{|u^+|} \right) \cdot \nu_u = 0 \quad (5.15)$$

$\mathcal{H}^1$ -a.e. on J_v .

From (5.15) and (5.14), we deduce that there exists $\lambda \in \mathbb{R}$ such that the following system of equations is satisfied $\mathcal{H}^1$ -a.e. on J_v :

$$\begin{cases} \frac{u_1^+}{|u^+|} - \frac{u_1^-}{|u^-|} = -\lambda(u_2^+ - u_2^-) \\ \frac{bu_2^+ - au_1^+}{|u^+|} - \frac{bu_2^- - au_1^-}{|u^-|} = \lambda(b(u_1^+ - u_1^-) + a(u_2^+ - u_2^-)). \end{cases}$$

By elementary linear algebra, using $b \neq 0$, we obtain the equivalent systems

$$\begin{cases} \frac{u_1^+}{|u^+|} - \frac{u_1^-}{|u^-|} = -\lambda(u_2^+ - u_2^-) \\ \frac{u_2^+}{|u^+|} - \frac{u_2^-}{|u^-|} = \lambda(u_1^+ - u_1^-), \end{cases} \iff \left(\frac{u^+}{|u^+|} - \frac{u^-}{|u^-|} \right) \cdot (u^+ - u^-) = 0,$$

that is $u^- \cdot u^+ = |u^-||u^+|$, $\mathcal{H}^1$ -a.e. on J_v . It follows that $u^- = \mu u^+$ $\mathcal{H}^1$ -a.e. on J_v for some $\mu : J_v \rightarrow (0, +\infty)$, and then $u^+ / |u^+| = u^- / |u^-|$ $\mathcal{H}^1$ -a.e. on J_v . This proves that $\mathcal{H}^1(J_v) = 0$ and thus $D^j v = 0$.

Now we prove that $D^c v = 0$. By assumption ii), the Cantor part of the measure

$$-\partial_1 u_2 + b\partial_2 u_1 + a\partial_2 u_2$$

vanishes. From $D^c u = \eta \otimes \xi |D^c u|$, we can compute $D_k^c u_l$ for $k, l = 1, 2$, and we find

$$(-\eta_2, b\eta_1 + a\eta_2) \cdot \xi = 0 \tag{5.16}$$

$|D^c u|$ -a.e. on Ω . By assumption iii), the Cantor part of the measure

$$(\partial_1 - a\partial_2)(u_1/|u|) + b\partial_2(u_2/|u|)$$

vanishes. Thus, letting $\vartheta = (\vartheta_1, \vartheta_2) = ((u^*)^\perp \otimes (u^*)^\perp)\eta$, we can use (5.4), (5.11) and (5.13) to compute $D_k^c(u_l/|u|)$ for $k, l = 1, 2$, and we find

$$(\vartheta_1, b\vartheta_2 - a\vartheta_1) \cdot \xi = 0 \tag{5.17}$$

$|D^c u|$ -a.e. on Ω . From (5.16) and (5.17), we deduce that there exists $\lambda \in \mathbb{R}$ such that

$$\begin{cases} \vartheta_1 = -\lambda\eta_2 \\ b\vartheta_2 - a\vartheta_1 = \lambda(b\eta_1 + a\eta_2) \end{cases} \iff \begin{cases} \vartheta_1 = -\lambda\eta_2 \\ \vartheta_2 = \lambda\eta_1. \end{cases}$$

Here we used $b \neq 0$. This, in turn, is equivalent with

$$\vartheta \cdot \eta = ((u^*)^\perp \otimes (u^*)^\perp)\eta \cdot \eta = 0$$

$|D^c u|$ -a.e. on Ω . Using the identity $((u^*)^\perp \otimes (u^*)^\perp)\eta \cdot \eta = ((u^*)^\perp \cdot \eta)^2$, we deduce that $(u^*)^\perp \cdot \eta = 0$, and thus $((u^*)^\perp \otimes (u^*)^\perp)\eta = 0$ as well. This proves that the expression $(\nabla F(u^*)\eta) \otimes \xi$ in (5.13) is equal to 0, whence $D^c v = 0$. $\square$

Theorem 5.1 has the following corollaries:

Corollary 5.2. *Let $C \subseteq \mathbb{H}$ be a convex isoperimetric set and let $f : D \rightarrow \mathbb{R}$ be the function in (4.1). Then we have*

$$\frac{\nabla f(z) + 2z^\perp}{|\nabla f(z) + 2z^\perp|} \in W_{\text{loc}}^{1,1}(\text{int}(D) \setminus \Sigma(f); \mathbb{R}^2). \tag{5.18}$$

Proof. The vector field $u(z) = (u_1(z), u_2(z)) = \nabla f(z) + 2z^\perp$ satisfies $\text{div } u^\perp = -4$ in $\text{int}(D)$. Moreover, by Proposition 4.1, we have $\text{div}(u/|u|) = H$ in $\text{int}(D) \setminus \Sigma(f)$. The claim follows from Theorem 5.1 with $a = 0$ and $b = 1$. $\square$

Corollary 5.3. *Let $C \subseteq \mathbb{H}$ be a convex isoperimetric set and let $h : E \rightarrow \mathbb{R}$ be the function in (4.7). Then we have*

$$\frac{u}{|u|} \in W_{\text{loc}}^{1,1}(\text{int}(E) \setminus (\{y = 0\} \cup \Sigma(h)); \mathbb{R}^2), \quad (5.19)$$

with $u = (u_1, u_2) = (h_y - 2hh_t, 1 - 2yh_t)$.

Proof. We use Theorem 5.1 with $a = 2h$ and $b = -2y$. The vector field u is in $BV_{\text{loc}}(\text{int}(E); \mathbb{R}^2)$ and satisfies

$$\mathcal{M}u^\perp = (\partial_y - 2h\partial_t)(2yh_t - 1) - 2y\partial_t(h_y - 2hh_t) = 2h_t(1 + 2yh_t) \in L_{\text{loc}}^\infty(\text{int}(E)).$$

Moreover, by Proposition 4.2, $u/|u|$ satisfies in distributional sense

$$\mathcal{M}\left(\frac{u}{|u|}\right) = (\partial_y - 2h\partial_t)\left(\frac{u_1}{|u|}\right) - 2y\partial_t\left(\frac{u_2}{|u|}\right) = H$$

in $\text{int}(E) \setminus \Sigma(h)$, where $H > 0$ is the curvature of C . We have

$$u/|u| \in BV_{\text{loc}}(\text{int}(E) \setminus \Sigma(h))$$

by Proposition 3.8. The claim follows from Theorem 5.1. $\square$

Observing that $2yh_y - 2h = 2y(h_y - 2hh_t) + 2h(2yh_t - 1)$, from Corollary 5.3 we also get the following

Corollary 5.4. *Let $C \subseteq \mathbb{H}$ be a convex isoperimetric set and let $h : E \rightarrow \mathbb{R}$ be the function in (4.7). Then*

$$\frac{(1 - 2yh_t, 2yh_y - 2h)}{|(h_y - 2hh_t, 1 - 2yh_t)|} \in W_{\text{loc}}^{1,1}(\text{int}(E) \setminus (\{y = 0\} \cup \Sigma(h)); \mathbb{R}^2). \quad (5.20)$$

We conclude this section with a remark concerning smooth regularizations of BV vector fields. Let $\chi \in C_c^\infty(\mathbb{R}^2)$ be a smoothing kernel with $\chi \geq 0$, $\chi(q) = 0$ if $|q| \geq 1$ and $\|\chi\|_1 = 1$. For $\varepsilon > 0$, let $\chi_\varepsilon(q) = \varepsilon^{-2}\chi(q/\varepsilon)$. Let Ω be an open subset of $\mathbb{R}^2$ and let $u \in L^1(\Omega; \mathbb{R}^2)$ be an integrable vector field. Then, for any q_0 in $\Omega_\varepsilon = \{q \in \Omega \mid \text{dist}(q; \partial\Omega) > \varepsilon\}$, denote by

$$u_\varepsilon(q_0) = \int_{\Omega} \chi_\varepsilon(q_0 - q)u(q) dq \quad (5.21)$$

the standard mollification of u at the point q_0 . If $u \in BV(\Omega; \mathbb{R}^2)$, then

$$\lim_{\varepsilon \downarrow 0} u_\varepsilon(q_0) = u^*(q_0) \quad \text{for } \mathcal{H}^1\text{-a.e. } q_0 \in \Omega,$$

where u^* is the precise representative of u defined in (5.3). In particular, if $q_0 \in J_u$, then $u^*(q_0) = \frac{1}{2}(u^+(q_0) + u^-(q_0))$, where $u^\pm(q_0) \in \mathbb{R}^2$ satisfy (5.2). We need the following variant of this property:

Lemma 5.5. *Let $u \in BV(\Omega; \mathbb{R}^2)$ and let $q_1 \in \mathbb{R}^2$ be a fixed vector with $|q_1| < 1$. Then we have*

$$\lim_{\varepsilon \downarrow 0} u_\varepsilon(q_0 + \varepsilon q_1) = u^*(q_0) \quad \text{for every } q_0 \in \Omega \setminus S_u. \quad (5.22)$$

Moreover, for all $q_0 \in J_u$, there exist numbers $\alpha, \beta \in [0, 1]$ (which may depend on q_0, q_1 and χ), such that $\alpha + \beta = 1$ and

$$\lim_{\varepsilon \downarrow 0} u_\varepsilon(q_0 + \varepsilon q_1) = \alpha u^+(q_0) + \beta u^-(q_0). \quad (5.23)$$

Proof. We prove only (5.23), the proof of (5.22) being analogous. We assume without loss of generality that the point $q_0 \in J_u$ is 0 and that the jump direction ν_u at 0 is $(0, 1)$. Let $u^+ = u^+(0)$ and $u^- = u^-(0)$ be vectors such that (5.2) holds at 0. Let $B_\varepsilon^\pm = \{q \in \mathbb{R}^2 \mid |\varepsilon q_1 - q| < \varepsilon, q \cdot (0, \pm 1) > 0\}$ for $\varepsilon > 0$ and let

$$\alpha = \int_{B_\varepsilon^+} \chi_\varepsilon(\varepsilon q_1 - q) dq, \quad \beta = \int_{B_\varepsilon^-} \chi_\varepsilon(\varepsilon q_1 - q) dq. \quad (5.24)$$

A change of variable shows that the definition of α and β does not depend on $\varepsilon > 0$. Using (5.24), we find

$$\begin{aligned} u_\varepsilon(\varepsilon q_1) &= \int_{B_\varepsilon^+} \chi_\varepsilon(\varepsilon q_1 - q)(u(q) - u^+) dq + \alpha u^+ \\ &\quad + \int_{B_\varepsilon^-} \chi_\varepsilon(\varepsilon q_1 - q)(u(q) - u^-) dq + \beta u^-. \end{aligned}$$

Now our claim (5.23) follows from (5.2) and from

$$\left| \int_{B_\varepsilon^\pm} \chi_\varepsilon(\varepsilon q_1 - q)(u(q) - u^\pm) dq \right| \leq c \|\chi\|_\infty \int_{B^\pm(0, 2\varepsilon)} |u(q) - u^\pm| dq.$$

□

6 Stability of the curvature

Let $\Omega \subseteq \mathbb{R}^2$ be a bounded open set, let $a, b \in \text{Lip}(\Omega)$ be Lipschitz functions and denote by $a_\varepsilon \in C^\infty(\Omega_\varepsilon)$ the usual mollification of a . We consider the differential operators $\mathcal{M}$ and $\mathcal{M}_\varepsilon$ defined by

$$\begin{aligned} \mathcal{M}u &= (\partial_1 - a\partial_2)u_1 + b\partial_2u_2 = \text{tr}(\nabla u B), \\ \mathcal{M}_\varepsilon u &= (\partial_1 - a_\varepsilon\partial_2)u_1 + b\partial_2u_2 = \text{tr}(\nabla u B_\varepsilon), \end{aligned} \quad (6.1)$$

where $u = (u_1, u_2)$,

$$B = \begin{pmatrix} 1 & 0 \\ -a & b \end{pmatrix}, \quad B_\varepsilon = \begin{pmatrix} 1 & 0 \\ -a_\varepsilon & b \end{pmatrix} \quad (6.2)$$

and tr denotes the trace of the 2×2 matrices $\nabla u B$ and $\nabla u B_\varepsilon$.

Let $f \in \text{Lip}(\Omega)$ be a Lipschitz function and denote by $f_\varepsilon \in C^\infty(\Omega_\varepsilon)$ the mollification of f . We define the vector fields

$$u = (\partial_1 f - a \partial_2 f, b \partial_2 f) + \omega \quad \text{and} \quad u_\varepsilon = (\partial_1 f_\varepsilon - a_\varepsilon \partial_2 f_\varepsilon, b \partial_2 f_\varepsilon) + \omega, \quad (6.3)$$

where $\omega \in \text{Lip}(\Omega; \mathbb{R}^2)$ is a Lipschitz continuous vector field.

Theorem 6.1 (Stability of the curvature). *Let $\Omega \subseteq \mathbb{R}^2$ be a bounded open set, let $a, b \in \text{Lip}(\Omega)$ with $b \neq 0$ in Ω , let $\omega \in \text{Lip}(\Omega; \mathbb{R}^2)$ and let $f \in \text{Lip}(\Omega)$ be a function such that $\nabla f \in BV(\Omega; \mathbb{R}^2)$. Let $u \in BV(\Omega; \mathbb{R}^2)$ and $u_\varepsilon \in BV(\Omega_\varepsilon; \mathbb{R}^2)$ be the vector fields defined in (6.3). Assume that:*

- i) *There exist $\delta > 0$ and $\varepsilon_0 > 0$ such that $|u(q)| \geq \delta$ for $\mathcal{L}^2$ -a.e. $q \in \Omega$ and $|u_\varepsilon(q)| \geq \delta$ for all $q \in \Omega_\varepsilon$, $0 < \varepsilon < \varepsilon_0$;*
- ii) *$\mathcal{M}u^\perp \in L^1(\Omega)$, where $u^\perp = (-u_2, u_1)$;*
- iii) *$\mathcal{M}\left(\frac{u}{|u|}\right) \in L^1(\Omega)$.*

Then

$$\lim_{\varepsilon \downarrow 0} \int_K \left| \mathcal{M}_\varepsilon\left(\frac{u_\varepsilon}{|u_\varepsilon|}\right) - \mathcal{M}\left(\frac{u}{|u|}\right) \right| dq = 0 \quad (6.4)$$

for any compact set $K \subseteq \Omega$.

Proof. Let $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a smooth mapping satisfying (5.10), let $v = F \circ u$ and $v_\varepsilon = F \circ u_\varepsilon$. By (6.1), assumption iii) and (5.7), we have

$$\mathcal{M}v = \text{tr}((\nabla F \circ u^*) \nabla u B), \quad \mathcal{M}_\varepsilon v_\varepsilon = \text{tr}((\nabla F \circ u_\varepsilon) \nabla u_\varepsilon B_\varepsilon), \quad (6.5)$$

where $\nabla u \in L^1(\Omega; M^{2 \times 2})$ is the density of the measure $D^a u$ with respect to Lebesgue measure. We denote by $\nabla w \in L^1(\Omega; M^{2 \times 2})$ the density with respect to Lebesgue measure of the absolutely continuous part of the distributional derivative of $w = \nabla f$. Letting $\vartheta = (-a, b)$, $\vartheta_\varepsilon = (-a_\varepsilon, b)$ and $w_\varepsilon = \nabla f_\varepsilon$, we have the relations

$$\begin{aligned} \nabla u &= B^T \nabla w + \Theta & \text{with } \Theta &= \partial_2 f \nabla \vartheta + \nabla \omega, \\ \nabla u_\varepsilon &= B_\varepsilon^T \nabla w_\varepsilon + \Theta_\varepsilon & \text{with } \Theta_\varepsilon &= \partial_2 f_\varepsilon \nabla \vartheta_\varepsilon + \nabla \omega. \end{aligned} \quad (6.6)$$

The gradient of w_ε is

$$\nabla w_\varepsilon(q) = \int_\Omega \chi_\varepsilon(q - q') dDw$$

for $q \in \Omega_\varepsilon$, where

$$Dw = D^a w + D^c w + D^j w = \nabla w \mathcal{L}^2 + D^c w + (w^+ - w^-) \otimes \nu_w \mathcal{H}^1 \llcorner J_w,$$

by the decomposition formulae (5.5)–(5.6). Thus we can split $\mathcal{M}_\varepsilon v_\varepsilon$ in the following way:

$$\mathcal{M}_\varepsilon v_\varepsilon = \text{tr}((\nabla F \circ u_\varepsilon) \Theta_\varepsilon B_\varepsilon) + \mathcal{M}_\varepsilon^a v_\varepsilon + \mathcal{M}_\varepsilon^c v_\varepsilon + \mathcal{M}_\varepsilon^j v_\varepsilon,$$

where

$$\begin{aligned}
\mathcal{M}_\varepsilon^a v_\varepsilon(q) &= \int_{\Omega} \chi_\varepsilon(q - q') \operatorname{tr} \left((\nabla F \circ u_\varepsilon) B_\varepsilon^T \nabla w(q') B_\varepsilon \right) dq', \\
\mathcal{M}_\varepsilon^c v_\varepsilon(q) &= \int_{\Omega} \chi_\varepsilon(q - q') \operatorname{tr} \left((\nabla F \circ u_\varepsilon) B_\varepsilon^T dD^c w(q') B_\varepsilon \right), \\
\mathcal{M}_\varepsilon^j v_\varepsilon(q) &= \int_{J_w} \chi_\varepsilon(q - q') \left((\nabla F \circ u_\varepsilon) B_\varepsilon^T (w^+(q') - w^-(q')) \right) \cdot (\nu_w(q') B_\varepsilon) d\mathcal{H}^1(q').
\end{aligned} \tag{6.7}$$

Here, unless specified otherwise, the functions depend on the variable q . To get the last line of (6.7), we also used the identities $A(\xi \otimes \eta) B_\varepsilon = (A\xi) \otimes (\eta^T B_\varepsilon)$ and $\operatorname{tr}(\xi \otimes \eta) = \xi \cdot \eta$.

Similarly, we have the decomposition

$$\mathcal{M}v = \operatorname{tr} \left((\nabla F \circ u^*) \Theta B \right) + \operatorname{tr} \left((\nabla F \circ u^*) B^T \nabla w B \right).$$

We claim that

$$\lim_{\varepsilon \downarrow 0} \int_K |\mathcal{M}_\varepsilon^j v_\varepsilon(q)| dq = 0. \tag{6.8}$$

There is a constant $c > 0$ such that for $q \in K$ and for all $0 < \varepsilon < \varepsilon_0$, we have $|B_\varepsilon(q)| \leq c$. By Minkowski's inequality and a change of variable, we find

$$\begin{aligned}
\int_K |\mathcal{M}_\varepsilon^j v_\varepsilon(q)| dq &\leq c \int_{J_w} \int_K \chi_\varepsilon(q - q') \left| \nabla F(u_\varepsilon(q)) B_\varepsilon^T(q) (w^+(q') - w^-(q')) \right| dq d\mathcal{H}^1 \\
&\leq c \|\chi\|_\infty \int_{J'_w} \int_{\{|q| < 1\}} \left| \nabla F(u_\varepsilon(q' + \varepsilon q)) B_\varepsilon^T(q' + \varepsilon q) (w^+(q') - w^-(q')) \right| dq d\mathcal{H}^1,
\end{aligned}$$

where J'_w is the intersection of J_w with a compact set $K' \subseteq \Omega$ which is independent of $0 < \varepsilon < \varepsilon_0$ (we choose a smaller ε_0 if necessary). Since the function in the last integral is uniformly bounded, we can invoke dominated convergence and pass to the limit $\varepsilon \downarrow 0$ inside the integral.

If $q' \in J_w$, then, by (5.23), we have

$$\lim_{\varepsilon \downarrow 0} w_\varepsilon(q' + \varepsilon q) = \alpha w^+(q') + \beta w^-(q'),$$

for suitable $\alpha, \beta \in [0, 1]$ which depend on q', q and χ . This implies

$$\lim_{\varepsilon \downarrow 0} \nabla F(u_\varepsilon(q' + \varepsilon q)) = \nabla F(\alpha u^+(q') + \beta u^-(q')),$$

where, according to (6.3) and (5.8), $u^\pm = (w_1^\pm - \alpha w_2^\pm, \beta w_2^\pm) + \omega$. On the other hand, by Theorem 5.1, we have $u/|u| \in W^{1,1}(\Omega; \mathbb{R}^2)$ and there exists a function $\mu : J_u \rightarrow (0, +\infty)$ such that $u^- = \mu u^+$ on J_u . Then the vectors $\alpha u^+(q') + \beta u^-(q')$ and $u^+(q') - u^-(q')$ are parallel. Now, from the identity

$$B^T(q')(w^+(q') - w^-(q')) = u^+(q') - u^-(q')$$

and from (5.11), we get

$$\nabla F(\alpha u^+(q') + \beta u^-(q')) B^T(q')(w^+(q') - w^-(q')) = 0$$

for all $q' \in J_w = J_u$. This finishes the proof of (6.8).

By dominated convergence, we find

$$\lim_{\varepsilon \downarrow 0} \int_K |\operatorname{tr}((\nabla F \circ u_\varepsilon) \Theta_\varepsilon B_\varepsilon - (\nabla F \circ u^*) \Theta B)| dq = 0. \quad (6.9)$$

With the short notation

$$L_\varepsilon(q, q') = |(\nabla F \circ u_\varepsilon) B_\varepsilon^T \nabla w(q') B_\varepsilon - (\nabla F \circ u^*) B^T \nabla w B|,$$

we have

$$\int_K |\mathcal{M}_\varepsilon^a v_\varepsilon - \operatorname{tr}((\nabla F \circ u^*) B^T \nabla w B)| dq \leq c \int_K \int_\Omega \chi_\varepsilon(q - q') L_\varepsilon(q, q') dq' dq.$$

As before, a function depends on the variable q unless stated otherwise. By the triangle inequality,

$$\begin{aligned} L_\varepsilon(q, q') &\leq |(\nabla F \circ u_\varepsilon) B_\varepsilon^T \nabla w(q') B_\varepsilon - (\nabla F \circ u_\varepsilon) B_\varepsilon^T \nabla w(q') B| \\ &\quad + |(\nabla F \circ u_\varepsilon) B_\varepsilon^T \nabla w(q') B - \nabla F(u^*(q')) B^T(q') \nabla w(q') B| \\ &\quad + |\nabla F(u^*(q')) B^T(q') \nabla w(q') B - (\nabla F \circ u^*) B^T \nabla w B|, \end{aligned}$$

and we get

$$\int_K |\mathcal{M}_\varepsilon^a v_\varepsilon - \operatorname{tr}((\nabla F \circ u^*) B^T \nabla w B)| dq \leq c \{I_\varepsilon + J_\varepsilon + K_\varepsilon\}, \quad (6.10)$$

where

$$\begin{aligned} I_\varepsilon &= \int_\Omega |\nabla w(q')| \int_K \chi_\varepsilon(q - q') |B_\varepsilon(q) - B(q)| dq dq', \\ J_\varepsilon &= \int_{K'} |\nabla w(q')| \int_{\{|q| < 1\}} |\nabla F(u_\varepsilon(q' + \varepsilon q)) B_\varepsilon^T(q' + \varepsilon q) - \nabla F(u^*(q')) B^T(q')| dq dq', \\ K_\varepsilon &= \int_{K'} \int_{\{|q| < 1\}} |[(\nabla F \circ u^*) B^T \nabla w](q') - [(\nabla F \circ u^*) B^T \nabla w](q' + \varepsilon q)| dq dq'. \end{aligned}$$

In the limit $\varepsilon \downarrow 0$, we have $I_\varepsilon \rightarrow 0$ by dominated convergence, $J_\varepsilon \rightarrow 0$ by dominated convergence and by (5.22) and $K_\varepsilon \rightarrow 0$ by the continuity in the mean for L^1 functions.

An analogous argument shows that

$$\lim_{\varepsilon \downarrow 0} \int_K |\mathcal{M}_\varepsilon^c v_\varepsilon(q)| dq = 0. \quad (6.11)$$

This concludes the proof of (6.4). $\square$

Remark 6.1. We use Theorem 6.1 in the following two situations.

1) In the coordinates $z = (x, y)$, we choose $a = 0$, $b = 1$ and $\omega(z) = 2z^\perp$. The vector field in (6.3) is $u(z) = \nabla f(z) + 2z^\perp$ and

$$\mathcal{M}\left(\frac{u}{|u|}\right) = \operatorname{div} \left(\frac{\nabla f(z) + 2z^\perp}{|\nabla f(z) + 2z^\perp|} \right)$$

is the curvature operator in (4.2).

2) In the coordinates $\zeta = (y, t)$, we have a function $h \in \operatorname{Lip}(\Omega)$ and we choose $a = 2h$, $b = -2y$, and $\omega = (0, 1)$. The vector field in (6.3) is $u = (u_1, u_2) = (h_y - 2hh_t, 1 - 2yh_t)$ – this is (4.9) – and

$$\mathcal{M}\left(\frac{u}{|u|}\right) = (\partial_y - 2h\partial_t)\left(\frac{u_1}{|u|}\right) - 2y\partial_t\left(\frac{u_2}{|u|}\right)$$

is the curvature operator in (4.8).

7 Foliation by geodesics

Let $u \in BV(\mathbb{R}^2; \mathbb{R}^2) \cap L^\infty(\mathbb{R}^2; \mathbb{R}^2)$ and, for $\varrho > 0$ and $q \in \mathbb{R}^2$, define

$$C_q([- \varrho, \varrho]; \mathbb{R}^2) = \left\{ \gamma \in C([- \varrho, \varrho]; \mathbb{R}^2) \mid \gamma(s) = q + \int_0^s u(\gamma(\sigma)) d\sigma, s \in [- \varrho, \varrho] \right\}.$$

Both $C([- \varrho, \varrho]; \mathbb{R}^2)$ and $C_q([- \varrho, \varrho]; \mathbb{R}^2)$ are endowed with the sup norm. Let $K \subseteq \mathbb{R}^2$ be a compact set. An $\mathcal{L}^2$ -measurable map $\Phi : K \rightarrow C([- \varrho, \varrho]; \mathbb{R}^2)$ is a Lagrangian flow starting from K relative to u if $\Phi(q) \in C_q([- \varrho, \varrho]; \mathbb{R}^2)$ for $\mathcal{L}^2$ -a.e. $q \in K$. By abuse of notation, we identify the map Φ with the map $\Phi : K \times [- \varrho, \varrho] \rightarrow \mathbb{R}^2$, $\Phi(q, s) = \Phi(q)(s)$. The flow is said to be regular if there exists a constant $m \geq 1$ such that

$$\frac{1}{m} \mathcal{L}^2(A) \leq \mathcal{L}^2(\Phi(A, s)) \leq m \mathcal{L}^2(A) \quad (7.12)$$

for all $\mathcal{L}^2$ -measurable sets $A \subseteq K$ and for all $s \in [- \varrho, \varrho]$.

Theorem 7.1 (Ambrosio). *Assume that:*

- i) $u \in BV(\mathbb{R}^2; \mathbb{R}^2) \cap L^\infty(\mathbb{R}^2; \mathbb{R}^2)$;
- ii) $\operatorname{div} u \in L^\infty(\mathbb{R}^2)$.

Then, for any compact set $K \subseteq \mathbb{R}^2$ and for all $\varrho > 0$, there exists a unique regular Lagrangian flow $\Phi : K \times [- \varrho, \varrho] \rightarrow \mathbb{R}^2$ starting from K relative to u , unique in the sense that any other regular Lagrangian flow for u starting from K coincides with Φ $\mathcal{L}^2$ -a.e. Moreover:

- 1) The flow is stable, in the sense that, if $\{u_\varepsilon\}_{0 < \varepsilon < \varepsilon_0}$ is a family of smooth vector fields which is uniformly bounded in $BV(\mathbb{R}^2; \mathbb{R}^2) \cap L^\infty(\mathbb{R}^2; \mathbb{R}^2)$ and such that the u_ε have uniformly bounded divergence, then $u_\varepsilon \rightarrow u$ in $L^1(\mathbb{R}^2; \mathbb{R}^2)$ implies

$$\lim_{\varepsilon \downarrow 0} \int_K \sup_{s \in [-\varrho, \varrho]} |\Phi_\varepsilon(q, s) - \Phi(q, s)| dq = 0, \quad (7.13)$$

where Φ_ε denotes the flow of u_ε . In particular, the mapping $(q, s) \mapsto \Phi(q, s)$ is $\mathcal{L}^2 \times \mathcal{L}^1$ -measurable.

- 2) The flow satisfies the semigroup property

$$\Phi(\Phi(q, s), \sigma) = \Phi(q, s + \sigma) \quad (7.14)$$

for all $s, \sigma \in [-\varrho, \varrho]$ with $s + \sigma \in [-\varrho, \varrho]$.

The existence statement is Theorem 6.2, the uniqueness statement is Theorem 6.4, the stability statement is Theorem 6.6, the semigroup property is Remark 6.7 in [A]. Ambrosio's theory holds more generally for non autonomous vector fields in any space dimension.

Remark 7.1.

- 1) If $\Omega \subseteq \mathbb{R}^2$ is an open set, $K \subseteq \Omega$ is compact and $u \in BV(\Omega; \mathbb{R}^2) \cap L^\infty(\Omega; \mathbb{R}^2)$ with $\operatorname{div} u \in L^\infty(\Omega)$, then there is a vector field in $BV(\mathbb{R}^2; \mathbb{R}^2) \cap L^\infty(\mathbb{R}^2; \mathbb{R}^2)$ with distributional divergence in $L^\infty(\mathbb{R}^2)$, which coincides with u in a neighbourhood of K . This vector field has a regular Lagrangian flow $\Phi : K \times [-\varrho, \varrho] \rightarrow \mathbb{R}^2$. If $\varrho > 0$ is sufficiently small, then, for almost every $q \in K$, $s \mapsto \Phi(q, s)$ is an integral curve of u and $\Phi(q, s) \in \Omega$ for all $s \in [-\varrho, \varrho]$.
- 2) If $N \subseteq \mathbb{R}^2$ is negligible, then $\Phi^{-1}(N) \subseteq K \times [-\varrho, \varrho]$ is also negligible. This follows from (7.12) by a Fubini-type argument.
- 3) If $F, G : \mathbb{R}^2 \rightarrow \mathbb{R}^k$ are $\mathcal{L}^2$ -measurable mappings such that $F = G$ $\mathcal{L}^2$ -a.e., then, for $\mathcal{L}^2$ -a.e. $q \in K$, $F(\Phi(q, s)) = G(\Phi(q, s))$ for a.e. $s \in [-\varrho, \varrho]$.
- 4) If $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ is locally Lipschitz continuous, then the chain rule holds along integral curves of u , i.e.

$$\frac{d}{ds} F(\Phi(q, s)) = \nabla F(\Phi(q, s)) \frac{d}{ds} \Phi(q, s) \quad (7.15)$$

for $\mathcal{L}^2$ -a.e. $q \in K$ and for a.e. $s \in [-\varrho, \varrho]$.

We are interested in the flow associated with a certain horizontal vector field tangent to the boundary of a convex isoperimetric set $C \subseteq \mathbb{H}$. Let $f : D \rightarrow \mathbb{R}$ be the convex function in (4.1). We consider the vector fields $v(z) = 2z - \nabla f^\perp(z)$ and $u(z) = v^\perp(z) = \nabla f(z) + 2z^\perp$ in $BV_{\operatorname{loc}}(\operatorname{int}(D); \mathbb{R}^2) \cap L^\infty_{\operatorname{loc}}(\operatorname{int}(D); \mathbb{R}^2)$. We have

$$\operatorname{div} v = 4 + f_{yx} - f_{xy} = 4. \quad (7.16)$$

The vector field v is the projection onto the xy -plane of the vector field

$$(x, y) \mapsto (f_y + 2x)\partial_x + (2y - f_x)\partial_y + (2xf_x + 2yf_y)\partial_t,$$

which is both horizontal and tangent to the graph of f at $\mathcal{H}^2$ -a.e. point.

Equation (4.2) completely determines the geometry of the integral curves of v .

Theorem 7.2 (Foliation by circles I). *Let $C \subseteq \mathbb{H}$ be a convex isoperimetric set with curvature $H > 0$ and $f : D \rightarrow \mathbb{R}$ be the function in (4.1). Let $K \subseteq \text{int}(D) \setminus \Sigma(f)$ be a compact set, $\Omega \Subset \text{int}(D) \setminus \Sigma(f)$ be an open neighbourhood of K and $\varrho > 0$ be sufficiently small. Denote by $\Phi : K \times [-\varrho, \varrho] \rightarrow \Omega$ the regular Lagrangian flow relative to the vector field $v(z) = 2z - \nabla f^\perp(z)$ starting from K . Then, for $\mathcal{L}^2$ -a.e. $z \in K$, the integral curve $s \mapsto \Phi(z, s)$ is an arc of circle with radius $1/H$ oriented clockwise.*

In the case of graphs of the type $x = h(y, t)$, the result is similar. Let $h : E \rightarrow \mathbb{R}$ be the function in (4.7). The vector field

$$v(\zeta) = (1 - 2yh_t, 2yh_y - 2h), \quad \zeta = (y, t) \in E, \quad (7.17)$$

is in $BV_{\text{loc}}(\text{int}(E); \mathbb{R}^2) \cap L_{\text{loc}}^\infty(\text{int}(E); \mathbb{R}^2)$, and

$$\text{div } v = -4h_t - 2yh_{ty} + 2yh_{yt} = -4h_t \in L_{\text{loc}}^\infty(\text{int}(E)). \quad (7.18)$$

The vector field v in (7.17) is the projection onto the yt -plane of the vector field

$$(y, t) \mapsto (h_y - 2hh_t)\partial_x + (1 - 2yh_t)\partial_y + (2yh_y - 2h)\partial_t, \quad (7.19)$$

which is both horizontal and tangent to the graph of h at $\mathcal{H}^2$ -a.e. point.

Theorem 7.3 (Foliation by circles II). *Let $C \subseteq \mathbb{H}$ be a convex isoperimetric set with curvature $H > 0$ and $h : E \rightarrow \mathbb{R}$ be the function in (4.7). Let $K \subseteq \text{int}(E) \setminus (\{y = 0\} \cup \Sigma(h))$ be a compact set, $\Omega \Subset \text{int}(E) \setminus (\{y = 0\} \cup \Sigma(h))$ be an open neighbourhood of K and $\varrho > 0$ be sufficiently small. Denote by $\Phi : K \times [-\varrho, \varrho] \rightarrow \Omega$ the regular Lagrangian flow relative to the vector field v in (7.17) starting from K . Then, for $\mathcal{L}^2$ -a.e. $\zeta \in K$, the projection onto the xy -plane of the curve*

$$s \mapsto (h(\Phi(\zeta, s)), \Phi(\zeta, s)) \in \mathbb{H}, \quad s \in [-\varrho, \varrho], \quad (7.20)$$

is an arc of circle with radius $1/H$ oriented clockwise.

We give two different proofs of Theorems 7.2 and 7.3 in Sections 8 and 9, respectively. The proof in Section 8 is based on Theorem 6.1 and on the stability property (7.13) of the flow, while the proof in Section 9 relies on a reparameterization argument involving (7.12). These theorems have the following geometric interpretation.

Corollary 7.4 (Foliation by geodesics I). *Under the assumptions of Theorem 7.2, for all $z \in K$, there is a geodesic $\gamma_z : [-\varrho, \varrho] \rightarrow \partial C$ with curvature H , such that $\gamma_z(0) = (z, f(z))$. Moreover, the length of γ_z is bounded from below by a positive constant depending on K .*

Proof. The horizontal lift of the plane curve $s \mapsto \Phi(z, s)$ given by Theorem 7.2, i.e. the curve

$$\gamma_z(s) = (\Phi(z, s), f(\Phi(z, s))), \quad s \in [-\varrho, \varrho],$$

is a geodesic with curvature H and with length bounded from below by a positive constant depending on K . By Theorem 7.2, this curves exist for $\mathcal{L}^2$ -a.e. $z \in K$. The stated result now follows from a density-compactness argument. $\square$

Analogously, we have:

Corollary 7.5 (Foliation by geodesics II). *Under the assumptions of Theorem 7.3, for all $\zeta \in K$, there is a geodesic $\gamma_\zeta : [-\varrho, \varrho] \rightarrow \partial C$ with curvature H , such that $\gamma_\zeta(0) = (h(\zeta), \zeta)$. Moreover, the length of γ_ζ is bounded from below by a positive constant depending on K .*

8 Approximation argument

We prove Theorems 7.2 and 7.3 using the stability property (7.13) of the flow and the stability property (6.4) of the curvature with respect to smooth approximations.

Proof of Theorem 7.2. For $\varepsilon > 0$ sufficiently small, the mollification f_ε of f is defined in some open neighbourhood of $\overline{\Omega}$. Let $v_\varepsilon = 2z - \nabla f_\varepsilon^\perp$. From (7.16), it follows that

$$\operatorname{div} v_\varepsilon = \operatorname{div} v = 4 \quad \text{in } \Omega. \quad (8.1)$$

In particular, $\operatorname{div} v_\varepsilon, \operatorname{div} v \in L^1(\Omega)$. Note that, by Proposition 3.7, there are constants $\delta > 0$ and $\varepsilon_0 > 0$, such that $|\nabla f_\varepsilon(z) + 2z^\perp| \geq \delta$ for all $z \in \Omega$ and $0 < \varepsilon < \varepsilon_0$, and $|\nabla f(z) + 2z^\perp| \geq \delta$ for $\mathcal{L}^2$ -a.e. $z \in \Omega$. Let $F \in C^\infty(\mathbb{R}^2; \mathbb{R}^2)$ be a mapping which satisfies (5.10). We define $w_\varepsilon = F \circ v_\varepsilon$ in Ω and $w = F \circ v$ in $\operatorname{int}(D)$. Then $w_\varepsilon = v_\varepsilon/|v_\varepsilon|$ and $w = v/|v|$ in Ω . Moreover,

$$\operatorname{div} w^\perp = H \quad (8.2)$$

in the weak sense, by Proposition 4.1 and Corollary 5.2.

Denote by $\Phi_\varepsilon : K \times [-\varrho, \varrho] \rightarrow \mathbb{R}^2$ the flow of v_ε . We have $\Phi_\varepsilon(K \times [-\varrho, \varrho]) \subseteq \Omega$ when ϱ is sufficiently small and $0 < \varepsilon < \varepsilon_0$. Denote by $J_\varepsilon(s) = J\Phi_\varepsilon(\cdot, s)$ the Jacobian in the z -variable of the flow. J_ε satisfies the differential equation $\dot{J}_\varepsilon = \operatorname{div} v_\varepsilon J_\varepsilon$, where the dot denotes the derivative with respect to the variable s . We find

$$J\Phi_\varepsilon(z, s) = e^{4s} \quad \text{for all } s \in [-\varrho, \varrho]. \quad (8.3)$$

We define the curvature function $H_\varepsilon : K \times [-\varrho, \varrho] \rightarrow \mathbb{R}$ to be

$$H_\varepsilon(z, s) = \operatorname{div} w_\varepsilon^\perp(\Phi_\varepsilon(z, s)) \quad \text{for all } z \in K \text{ and } s \in [-\varrho, \varrho]. \quad (8.4)$$

For $z \in K$, the integral curve $s \mapsto \Phi_\varepsilon(z, s)$ is of class C^∞ . The unit tangent vector $\Gamma_\varepsilon(z, s) = F(\dot{\Phi}_\varepsilon(z, s)) = w_\varepsilon(\Phi_\varepsilon(z, s))$ satisfies the equation

$$\dot{\Gamma}_\varepsilon(z, s) = -H_\varepsilon(z, s)\lambda_\varepsilon(z, s)\Gamma_\varepsilon^\perp(z, s) \quad \text{for all } z \in K \text{ and } s \in [-\varrho, \varrho], \quad (8.5)$$

where $\lambda_\varepsilon(z, s) = |v_\varepsilon(\Phi_\varepsilon(z, s))|$. From (8.5), we obtain

$$\int_K \psi \int_{-\varrho}^{\varrho} \left\{ \Gamma_\varepsilon \cdot \dot{\varphi} - H_\varepsilon \lambda_\varepsilon \Gamma_\varepsilon^\perp \cdot \varphi \right\} ds dz = 0 \quad (8.6)$$

for arbitrary test functions $\varphi \in C_c^1((-\varrho, \varrho); \mathbb{R}^2)$ and $\psi \in C_c^1(K)$. Our aim is to prove that

$$\int_K \psi \int_{-\varrho}^{\varrho} \left\{ \Gamma \cdot \dot{\varphi} - H \lambda \Gamma^\perp \cdot \varphi \right\} ds dz = 0, \quad (8.7)$$

with $\Gamma(z, s) = w(\Phi(z, s))$ and $\lambda(z, s) = |v(\Phi(z, s))|$.

We need some preliminary observations. By the stability property (7.13) of the Lagrangian flow, there is a sequence $\{\varepsilon_k\}_{k \in \mathbb{N}}$ such that $\varepsilon_k \downarrow 0$ and $\Phi_{\varepsilon_k} \rightarrow \Phi$ pointwise almost everywhere in $K \times [-\varrho, \varrho]$. We claim that

$$\lim_{k \rightarrow \infty} \int_K \int_{-\varrho}^{\varrho} |w_{\varepsilon_k} \circ \Phi_{\varepsilon_k} - w \circ \Phi| ds dz = 0. \quad (8.8)$$

Indeed, by (8.3), we have

$$\int_K |w_\varepsilon \circ \Phi_\varepsilon - w \circ \Phi_\varepsilon| dz \leq e^{-4s} \int_\Omega |w_\varepsilon - w| dz, \quad (8.9)$$

and the right hand side of (8.9) goes to zero. Moreover, we have

$$\lim_{k \rightarrow \infty} \int_K \int_{-\varrho}^{\varrho} |w \circ \Phi_{\varepsilon_k} - w \circ \Phi| ds dz = 0. \quad (8.10)$$

Indeed, let $\eta > 0$ be a positive number and choose a vector field $\tilde{w} \in C_c(\Omega; \mathbb{R}^2)$ such that $\|w - \tilde{w}\|_{L^1(\Omega; \mathbb{R}^2)} \leq \eta$. Denote by $\|\cdot\|$ the norm in $L^1(K \times [-\varrho, \varrho]; \mathbb{R}^2)$. By (7.12) and (8.3), we have

$$\begin{aligned} \|w \circ \Phi_\varepsilon - w \circ \Phi\| &\leq \|w \circ \Phi_\varepsilon - \tilde{w} \circ \Phi_\varepsilon\| + \|\tilde{w} \circ \Phi_\varepsilon - \tilde{w} \circ \Phi\| + \|\tilde{w} \circ \Phi - w \circ \Phi\| \\ &\leq c\eta + \|\tilde{w} \circ \Phi_\varepsilon - \tilde{w} \circ \Phi\|, \end{aligned}$$

where $c > 0$ does not depend on ε . This finishes the proof of (8.8). From (8.8), it follows that

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_K \psi \int_{-\varrho}^{\varrho} H \lambda \Gamma_{\varepsilon_k}^\perp \cdot \varphi ds dz &= \int_K \psi \int_{-\varrho}^{\varrho} H \lambda \Gamma^\perp \cdot \varphi ds dz, \\ \lim_{k \rightarrow \infty} \int_K \psi \int_{-\varrho}^{\varrho} \Gamma_{\varepsilon_k} \cdot \dot{\varphi} ds dz &= \int_K \psi \int_{-\varrho}^{\varrho} \Gamma \cdot \dot{\varphi} ds dz. \end{aligned} \quad (8.11)$$

Finally, performing the change of variable $z \mapsto \Phi_\varepsilon(z, -s)$ with area factor given by (8.3), we obtain

$$\int_K |(\operatorname{div} w_\varepsilon^\perp) \circ \Phi_\varepsilon - H| dz \leq e^{-4s} \int_\Omega |\operatorname{div} w_\varepsilon^\perp - H| dz, \quad (8.12)$$

for any $s \in [-\varrho, \varrho]$. Due to (6.4) in Theorem 6.1 with $a = 0$, $b = 1$ and $\omega = 2z^\perp$, the right hand side of (8.12) converges to 0 as $\varepsilon \downarrow 0$ (we have $\mathcal{M} = \operatorname{div}$). From (8.12), we deduce

$$\lim_{k \rightarrow \infty} \int_K \psi \int_{-\varrho}^{\varrho} (H_{\varepsilon_k} \lambda_{\varepsilon_k} - H \lambda) \Gamma_{\varepsilon_k}^\perp \cdot \varphi ds dz = 0. \quad (8.13)$$

Now, (8.7) follows from (8.11) and (8.13).

Since $\psi \in C_c^1(K)$ is arbitrary in (8.7), we deduce that the equation

$$\int_{-\varrho}^{\varrho} \left\{ \Gamma(z, s) \cdot \dot{\varphi}(s) - H\lambda(z, s)\Gamma^\perp(z, s) \cdot \varphi(s) \right\} ds = 0, \quad (8.14)$$

holds for $\mathcal{L}^2$ -a.e. $z \in K$ and for all $\varphi \in C_c^1((-\varrho, \varrho); \mathbb{R}^2)$. Now, by a standard argument, we conclude that for $\mathcal{L}^2$ -a.e. $z \in K$ we have

$$\dot{\Gamma}(z, s) = -H\lambda(z, s)\Gamma^\perp(z, s)$$

for a.e. $s \in [-\varrho, \varrho]$, where $\lambda(z, t) = |v(\Phi(z, t))| = |\dot{\Phi}(z, t)|$ is the length factor of the curve $s \mapsto \Phi(z, s)$. Then $\Phi(z, \cdot)$ parameterizes an arc of circle with curvature H . $\square$

Proof of Theorem 7.3. For $\varepsilon > 0$ sufficiently small, the mollification h_ε of h is defined in some open neighbourhood of $\bar{\Omega}$. We consider the vector fields

$$v_\varepsilon = (1 - 2y\partial_t h_\varepsilon, 2y\partial_y h_\varepsilon - 2h_\varepsilon) \quad \text{and} \quad u_\varepsilon = (\partial_y h_\varepsilon - 2h_\varepsilon \partial_t h_\varepsilon, 1 - 2y\partial_t h_\varepsilon).$$

Analogously, in $\text{int}(E)$, we can define

$$v = (1 - 2yh_t, 2yh_y - 2h) \quad \text{and} \quad u = (h_y - 2hh_t, 1 - 2yh_t).$$

By Proposition 3.8, there are constants $\delta > 0$ and $\varepsilon_0 > 0$, such that $|u_\varepsilon| \geq \delta$ in Ω for $0 < \varepsilon < \varepsilon_0$, and $|u| \geq \delta$ $\mathcal{L}^2$ -a.e. in Ω . Let $F \in C^\infty(\mathbb{R}^2; \mathbb{R}^2)$ be a mapping which satisfies (5.10). We can define the vector fields $w_\varepsilon = F \circ u_\varepsilon$ in Ω and $w = F \circ u$ in $\text{int}(E)$. Then $w_\varepsilon = u_\varepsilon/|u_\varepsilon|$ and $w = u/|u|$ in Ω .

Let $\mathcal{M}$ be the differential operator in (5.9), with $a = 2h$ and $b = -2y$. By Proposition 4.2 and Corollary 5.3, the vector field u satisfies

$$\mathcal{M}\left(\frac{u}{|u|}\right) = H \quad \text{in } \Omega,$$

in the weak sense. A direct computation shows that

$$\mathcal{M}u^\perp = 2h_t(1 + 2yh_t) \in L^\infty(\Omega).$$

Moreover, from (7.18), we get

$$\text{div } v_\varepsilon = -4\partial_t h_\varepsilon \quad \text{in } \Omega,$$

and there is a constant $c > 0$ such that $|\text{div } v_\varepsilon| \leq c$ in Ω for all $0 < \varepsilon < \varepsilon_0$. Let us denote by $\Phi_\varepsilon : K \times [-\varrho, \varrho] \rightarrow \mathbb{R}^2$ the flow of v_ε . As in the proof of Theorem 7.2, ϱ can be chosen in such a way that $\Phi_\varepsilon(K \times [-\varrho, \varrho]) \subseteq \Omega$ for $0 < \varepsilon < \varepsilon_0$, and the Jacobian of Φ_ε satisfies

$$c_1 \leq J\Phi_\varepsilon(\zeta, s) \leq c_2 \quad \text{for all } \zeta \in K \text{ and } s \in [-\varrho, \varrho], \quad (8.15)$$

with constants $0 < c_1 \leq c_2$ which do not depend on ε .

Let $s \mapsto \Gamma_\varepsilon(\zeta, s) = w_\varepsilon(\Phi_\varepsilon(\zeta, s))$ be the unit tangent vector of the projection onto the xy -plane of the curve $s \mapsto (h_\varepsilon(\Phi_\varepsilon(\zeta, s)), \Phi_\varepsilon(\zeta, s)) \in \mathbb{H}$. This tangent vector satisfies the equation

$$\dot{\Gamma}_\varepsilon(\zeta, s) = -H_\varepsilon(\zeta, s)\lambda_\varepsilon(\zeta, s)\Gamma_\varepsilon^\perp(\zeta, s) \quad \text{for all } \zeta \in K \text{ and } s \in [-\varrho, \varrho], \quad (8.16)$$

where $H_\varepsilon : K \times [-\varrho, \varrho] \rightarrow \mathbb{R}$ is the curvature function

$$H_\varepsilon(\zeta, s) = \mathcal{M}\left(\frac{u_\varepsilon}{|u_\varepsilon|}\right)(\Phi_\varepsilon(\zeta, s)), \quad (8.17)$$

and $\lambda_\varepsilon(\zeta, s) = |u_\varepsilon(\Phi_\varepsilon(\zeta, s))|$. The proof of formula (8.16) is postponed to Section 9, where it is proved under weaker regularity assumptions (see the proof just before formula (9.9)).

As in (8.6), we arrive at the integral equation

$$\int_K \psi \int_{-\varrho}^{\varrho} \left\{ \Gamma_\varepsilon \cdot \dot{\varphi} - H_\varepsilon \lambda_\varepsilon \Gamma_\varepsilon^\perp \cdot \varphi \right\} ds d\zeta = 0, \quad (8.18)$$

with $\varphi \in C_c^1((-\varrho, \varrho); \mathbb{R}^2)$ and $\psi \in C_c^1(K)$. Now the proof runs along the same lines as the proof of Theorem 7.2 above, and we can omit the details. Theorem 6.1 is used with $a = 2h$, $b = -2y$ and $\omega = (0, 1)$. $\square$

9 Reparameterization argument

Let $\Omega \subseteq \mathbb{R}^2$ be an open set and let v be a vector field such that

$$v \in BV(\Omega; \mathbb{R}^2) \cap L^\infty(\Omega; \mathbb{R}^2) \quad \text{and} \quad \operatorname{div} v \in L^\infty(\Omega). \quad (9.1)$$

The function v is defined pointwise, i.e. we choose a representative in the equivalence class of v . Our results hold independently of this choice. However, there is an exceptional set of points which may a priori depend on the representative.

Given a compact set $K \subseteq \Omega$ and a sufficiently small $\varrho > 0$, there exists a unique regular Lagrangian flow $\Phi : K \times [-\varrho, \varrho] \rightarrow \Omega$ starting from K relative to v (Theorem 7.1). Let $\lambda : \Omega \rightarrow \mathbb{R}$ be a measurable function such that

$$0 < c_1 \leq \lambda \leq c_2 \quad \mathcal{L}^2\text{-a.e. in } \Omega. \quad (9.2)$$

Then, for $\mathcal{L}^2$ -a.e. $q \in K$, the curve $s \mapsto \lambda(\Phi(q, s))$ is measurable and

$$c_1 \leq \lambda(\Phi(q, s)) \leq c_2 \quad \text{for a.e. } s \in [-\varrho, \varrho].$$

Thus, for $\mathcal{L}^2$ -a.e. $q \in K$, the change of parameter $\sigma_q : [-\varrho, \varrho] \rightarrow [\sigma_q(-\varrho), \sigma_q(\varrho)]$ defined by

$$\sigma_q(s) = \int_0^s \lambda(\Phi(q, \xi)) d\xi$$

is bi-Lipschitz, strictly increasing and admits therefore a bi-Lipschitz, strictly increasing inverse $\tau_q : [\sigma_q(-\varrho), \sigma_q(\varrho)] \rightarrow [-\varrho, \varrho]$, which satisfies

$$\dot{\tau}_q(s) = \frac{1}{\dot{\sigma}_q(\tau_q(s))} = \frac{1}{\lambda(\Phi(q, \tau_q(s)))} \quad \text{for a.e. } s \in [\sigma_q(-\varrho), \sigma_q(\varrho)]. \quad (9.3)$$

Consequently, for $\mathcal{L}^2$ -a.e. $q \in K$, the curve $\gamma_q : [\sigma_q(-\varrho), \sigma_q(\varrho)] \rightarrow \Omega$ defined by

$$\gamma_q(s) = \Phi(q, \tau_q(s)) \quad (9.4)$$

is absolutely continuous and satisfies

$$\dot{\gamma}_q(s) = \frac{v(\gamma_q(s))}{\lambda(\gamma_q(s))} \quad \text{for a.e. } s \in [\sigma_q(-\varrho), \sigma_q(\varrho)], \quad (9.5)$$

i.e. γ_q is an integral curve of the vector field v/λ .

Theorem 9.1 (Chain rule for integral curves). *Let $\Omega \subseteq \mathbb{R}^2$ be an open set, let $K \subseteq \Omega$ be a compact subset, let $w \in W^{1,1}(\Omega; \mathbb{R}^2)$ and let $\varrho > 0$ be sufficiently small. Then, for $\mathcal{L}^2$ -a.e. $q \in K$, the curve $w \circ \gamma_q$ with γ_q defined in (9.4) belongs to $W^{1,1}((\sigma_q(-\varrho), \sigma_q(\varrho)); \mathbb{R}^2)$, and its weak derivative is $(\nabla w \circ \gamma_q)\dot{\gamma}_q$.*

Proof. Let $\{w_k\}_{k \in \mathbb{N}}$ be a sequence of smooth vector fields $w_k \in W^{1,1}(\Omega; \mathbb{R}^2)$ converging to w in $W^{1,1}(\Omega; \mathbb{R}^2)$. Notice that the map $(q, s) \mapsto w(\Phi(q, s))$ is measurable and integrable. By Fubini's theorem and by the bounded volume distortion property (7.12) of the flow, we have

$$\int_K \int_{-\varrho}^{\varrho} |w_k(\Phi(q, s)) - w(\Phi(q, s))| ds dq \leq m \int_{-\varrho}^{\varrho} \int_{\Phi(K, s)} |w_k(q) - w(q)| dq ds,$$

and

$$\begin{aligned} \int_K \int_{-\varrho}^{\varrho} |(\nabla w_k(\Phi(q, s)) - \nabla w(\Phi(q, s)))v(\Phi(q, s))| ds dq &\leq \\ &\leq \|v\|_{\infty} \int_{-\varrho}^{\varrho} \int_K |\nabla w_k(\Phi(q, s)) - \nabla w(\Phi(q, s))| dq ds \\ &\leq m \|v\|_{\infty} \int_{-\varrho}^{\varrho} \int_{\Phi(K, s)} |\nabla w_k(q) - \nabla w(q)| dq ds. \end{aligned}$$

Notice that the curve $s \mapsto w_k(\Phi(q, s))$ is Lipschitz, for all $k \in \mathbb{N}$ and for $\mathcal{L}^2$ -a.e. $q \in K$, with derivative $s \mapsto \nabla w_k(\Phi(q, s))v(\Phi(q, s))$. Passing to a subsequence and relabelling if necessary, we conclude that

$$\lim_{k \rightarrow \infty} w_k(\Phi(q, \cdot)) = w(\Phi(q, \cdot)) \quad \text{in } W^{1,1}((-\varrho, \varrho); \mathbb{R}^2)$$

for $\mathcal{L}^2$ -a.e. $q \in K$, and that the weak derivative of $w(\Phi(q, \cdot))$ is $\nabla w(\Phi(q, \cdot))v(\Phi(q, \cdot))$.

Let $\varphi \in C_c^\infty((\sigma_q(-\varrho), \sigma_q(\varrho)); \mathbb{R}^2)$. Combining the previous observation, (9.3) and (9.5), we get

$$\begin{aligned} \int_{\sigma_q(-\varrho)}^{\sigma_q(\varrho)} w(\gamma_q(s)) \dot{\varphi}(s) ds &= \int_{-\varrho}^{\varrho} w(\Phi(q, s)) \dot{\varphi}(\sigma_q(s)) \dot{\sigma}_q(s) ds \\ &= - \int_{-\varrho}^{\varrho} \nabla w(\Phi(q, s)) v(\Phi(q, s)) \varphi(\sigma_q(s)) ds \\ &= - \int_{\sigma_q(-\varrho)}^{\sigma_q(\varrho)} \nabla w(\gamma_q(s)) \dot{\gamma}_q(s) \varphi(s) ds. \end{aligned}$$

Hence $w \circ \gamma_q \in W^{1,1}((\sigma_q(-\varrho), \sigma_q(\varrho)); \mathbb{R}^2)$, and its weak derivative is $(\nabla w \circ \gamma_q) \dot{\gamma}_q$. $\square$

By means of Theorem 9.1, we can now prove Theorems 7.2 and 7.3.

Proof of Theorem 7.2. We have $v(z) = 2z - \nabla f^\perp(z) \in BV(\Omega; \mathbb{R}^2) \cap L^\infty(\Omega; \mathbb{R}^2)$ and $\operatorname{div} v = 4$ (cf. (7.16)). The function $\lambda : \Omega \rightarrow \mathbb{R}$, $\lambda = |v|$, satisfies $0 < c_1 \leq \lambda \leq c_2$ $\mathcal{L}^2$ -a.e. in Ω by (3.6). By Corollary 5.2, the vector field $w = v/\lambda$ belongs to $W^{1,1}(\Omega; \mathbb{R}^2)$, and $\operatorname{div} w^\perp = H$ $\mathcal{L}^2$ -a.e. in Ω by (4.2).

Let $\gamma_z : (\sigma_z(-\varrho), \sigma_z(\varrho)) \rightarrow \Omega$, be the integral curve of the vector field w defined in (9.4) and satisfying (9.5). We claim that γ_z parameterizes an arc of circle with curvature H . Since $\Phi(z, \cdot)$ is a reparameterization of γ_z , proving the claim concludes the proof of Theorem 7.2.

By Remark 7.1, the following identities hold a.e. in $(\sigma_z(-\varrho), \sigma_z(\varrho))$ for $\mathcal{L}^2$ -a.e. $z \in K$:

- i) $|w \circ \gamma_z| = 1$,
- ii) $(w_1 \partial_x w_1 + w_2 \partial_x w_2) \circ \gamma_z = (w_1 \partial_y w_1 + w_2 \partial_y w_2) \circ \gamma_z = 0$ and
- iii) $(\operatorname{div} w^\perp) \circ \gamma_z = H$.

By Theorem 9.1, we find that $\gamma_z \in W^{2,1}((\sigma_z(-\varrho), \sigma_z(\varrho)); \mathbb{R}^2)$ with

$$\ddot{\gamma}_z = (\nabla w \circ \gamma_z) \dot{\gamma}_z. \quad (9.6)$$

Using (9.6), ii) and iii), we compute

$$\ddot{\gamma}_z \cdot \dot{\gamma}_z^\perp = (-\operatorname{div} w^\perp) \circ \gamma_z = -H.$$

Moreover, by i), we also have $\ddot{\gamma}_z \cdot \dot{\gamma}_z = 0$ a.e. in $(\sigma_z(-\varrho), \sigma_z(\varrho))$. Then $\ddot{\gamma}_z = -H \dot{\gamma}_z^\perp$ a.e. and this implies that γ_z is of class C^∞ . The claim follows. $\square$

Proof of Theorem 7.3. The vector field v in (7.17) is in $BV(\Omega; \mathbb{R}^2) \cap L^\infty(\Omega; \mathbb{R}^2)$ and satisfies $\operatorname{div} v = -4h_t \in L^\infty(\Omega)$ (cf. (7.18)).

We denote by $u = (h_y - 2hh_t, 1 - 2yh_t)$ the projection of the vector field (7.19) onto the xy -plane. The relation between v and u is

$$v = \begin{pmatrix} 0 & 1 \\ 2y & -2h \end{pmatrix} u. \quad (9.7)$$

In (9.7), we think of v and u as column vectors. The function $\lambda : \Omega \rightarrow \mathbb{R}$, $\lambda = |u|$, satisfies $0 < c_1 \leq \lambda \leq c_2$ $\mathcal{L}^2$ -a.e. in Ω by (3.9). Let $F \in C^\infty(\mathbb{R}^2; \mathbb{R}^2)$ be a mapping which satisfies (5.10) (with c_1 in place of δ). We consider the vector fields v/λ and $w = F \circ u$ in Ω . Then $w = u/|u|$ a.e. in Ω . We have $w \in W^{1,1}(\Omega; \mathbb{R}^2)$ by Corollary 5.3 and $v/\lambda \in W^{1,1}(\Omega; \mathbb{R}^2)$ by Corollary 5.4.

Denote by $\gamma_\zeta : [\sigma_\zeta(-\varrho), \sigma_\zeta(\varrho)] \rightarrow \Omega$ the integral curve of v/λ defined in (9.4). Then the curve $s \mapsto w(\gamma_\zeta(s))$ belongs to $W^{1,1}((\sigma_\zeta(-\varrho), \sigma_\zeta(\varrho)); \mathbb{R}^2)$ for $\mathcal{L}^2$ -a.e. $\zeta \in K$, and its weak derivative is equal to $(\nabla w \circ \gamma_\zeta) \dot{\gamma}_\zeta$ (Theorem 9.1). We claim that

$$\nabla w \frac{v}{\lambda} = -Hw^\perp \quad \mathcal{L}^2\text{-a.e. in } \Omega. \quad (9.8)$$

Indeed, using (5.11), we compute

$$\begin{aligned} \nabla w \frac{v}{\lambda} &= (\nabla F \circ u) \nabla u \frac{v}{\lambda} = \frac{1}{|u|^4} (u^\perp \otimes u^\perp) \nabla u v \\ &= \frac{1}{|u|^3} (u^\perp \cdot (\nabla u v)) w^\perp. \end{aligned}$$

On the other hand, from (4.8), we have $\mathcal{M}w = H$ in Ω , where $\mathcal{M}$ is the differential operator appearing in (5.9) with $a = 2h$ and $b = -2y$. Using (5.11) and the notation from (6.1), we get

$$\begin{aligned} \mathcal{M}w &= \operatorname{tr}((\nabla F \circ u) \nabla u B) = \frac{1}{|u|^3} \operatorname{tr}((u^\perp \otimes u^\perp) \nabla u B) \\ &= \frac{1}{|u|^3} u^\perp \cdot (u^\perp \nabla u B). \end{aligned}$$

Using (9.7), by a short computation, we find

$$u^\perp \cdot (u^\perp \nabla u B) = -u^\perp \cdot (\nabla u v),$$

and this ends the proof of (9.8). From Remark 7.1, it follows that the identity (9.8) holds along $\mathcal{L}^2$ -a.e. curve γ_ζ . Hence, by Theorem 9.1 applied to w , we have

$$\frac{d}{ds}(w \circ \gamma_\zeta) = -H(w^\perp \circ \gamma_\zeta) \quad (9.9)$$

in the sense of weak derivatives.

The projection of the vector field $\zeta = (y, t) \mapsto (h(\zeta), \zeta)$ onto the xy -plane belongs to $W^{1,1}(\Omega; \mathbb{R}^2)$. Denote by $\kappa_\zeta : [\sigma_\zeta(-\varrho), \sigma_\zeta(\varrho)] \rightarrow \mathbb{R}^2$ the projection of the

curve $(h(\gamma_\zeta), \gamma_\zeta)$ onto the xy -plane. This projection is a reparameterization of the curve in (7.20). We claim that κ_ζ parameterizes an arc of circle with curvature H oriented clockwise. By Theorem 9.1, for $\mathcal{L}^2$ -a.e. $\zeta \in K$, we have $\kappa_\zeta \in W^{1,1}((\sigma_\zeta(-\varrho), \sigma_\zeta(\varrho)); \mathbb{R}^2)$, and a short computation shows that the weak derivative of κ_ζ is $\dot{\kappa}_\zeta = w \circ \gamma_\zeta$. From (9.9), we deduce $\kappa_\zeta \in W^{2,1}((\sigma_\zeta(-\varrho), \sigma_\zeta(\varrho)); \mathbb{R}^2)$ and $\ddot{\kappa}_\zeta = -H\dot{\kappa}_\zeta^\perp$. This implies that κ_ζ is of class C^∞ and the claim follows. $\square$

10 Characterization of convex isoperimetric sets

In this final section, we combine the geometric description of the characteristic set of a convex set (Theorem 3.5) with the “foliation by geodesics property” of the boundary of convex isoperimetric sets (Corollaries 7.4 and 7.5), in order to prove the Characterization Theorem 1.1.

Proof of Theorem 1.1. According to Theorem 3.5, the characteristic set $\Sigma(C)$ of a convex isoperimetric set C admits a disjoint decomposition $\Sigma(C) = \Sigma^- \cup \Sigma^+$, where Σ^- and Σ^+ are two closed, horizontal segments. The curvature H of C scales like $1/\lambda$ with respect to the dilations δ_λ , $\lambda > 0$. Thus, we can assume $H = 2$ without loss of generality.

Claim 1. For all $p \in \partial C \setminus \Sigma(C)$, there is a curve of maximal length passing through p and contained in ∂C whose projection onto the xy -plane is an arc of circle with curvature H . One endpoint of the curve belongs to Σ^- , the other to Σ^+ .

Let $f : D \rightarrow \mathbb{R}$ be the function in (4.1), and let $h : E \rightarrow \mathbb{R}$ be the function in (4.7). Using an isometry and a left translation in the y direction if necessary, we can assume that any point $p \in \partial C \setminus \Sigma(C)$ is either of the form

- 1) $p = (z, f(z))$ for some $z \in \text{int}(D) \setminus \Sigma(f)$, or of the form
- 2) $p = (h(\zeta), \zeta)$ for some $\zeta \in \text{int}(E) \setminus (\{y = 0\} \cup \Sigma(h))$.

Then the existence of a piece of geodesic with curvature H passing through p and contained in ∂C follows either from Corollary 7.4 or from Corollary 7.5. The existence of a curve with maximal length whose projection onto the xy -plane is an arc of circle with curvature H follows from a compactness argument. If one endpoint of the maximal curve does not belong to $\Sigma(C)$, then a continuation argument involving the Uniqueness Lemma 3.9 provides a proper extension of the curve, which is impossible. Finally, the endpoints of the maximal curve cannot both belong to Σ^- or Σ^+ . This follows from our discussion of geodesics in Section 2.

Claim 2. The closed horizontal segments Σ^- and Σ^+ are points.

Suppose by contradiction that Σ^- is a closed, horizontal segment of positive length. After an isometry and a left translation, we have $\Sigma^- = \{(0, y, 0) \in \mathbb{H} \mid |y| \leq y_0\}$

for some $y_0 > 0$. Since the horizontal planes H_{p_0} and H_{-p_0} with $p_0 = (0, y_0, 0)$ are supporting planes for C , we have

$$C \subseteq H_{p_0}^+ \cap H_{-p_0}^+. \quad (10.1)$$

Claim 1 combined with an approximation-compactness argument shows that there is a geodesic $\gamma : [0, L] \rightarrow \partial C$ parameterized proportionally to arc-length, with curvature $H = 2$ and such that $\gamma(0) = 0$. We have $\dot{\gamma}(0) = (\alpha, \beta, 0)$ for some $\alpha, \beta \in \mathbb{R}$ with $|(\alpha, \beta)| = 1/2$. Assume $\alpha = 0$ and $\beta = 1/2$ (if $\alpha \neq 0$, the proof is easier). Then, by (2.4),

$$\gamma(s) = \frac{1}{2}(1 - \cos s, \sin s, s - \sin s) \in \partial C \quad \text{for all } s \in [0, L].$$

However, for any $c_0 > 0$, there is $\delta > 0$ with

$$s - \sin s < c_0(1 - \cos s) \quad \text{for all } s \in (0, \delta).$$

This contradicts (10.1).

Claim 3. Up to a left translation, the set C is of the form (1.8).

After a left translation, we can assume that $\Sigma^- = \{(0, -\pi/2)\}$ and that $\Sigma^+ = \{(z, t)\}$ with $t > -\pi/2$. This forces $z = 0$ and $t = \pi/2$, otherwise there would be at most one geodesic with curvature $H = 2$ connecting Σ^- to Σ^+ and, by Claim 1, this is not possible.

We claim that $0 \in \text{int}(D)$. If, by contradiction, $0 \in \partial D$, then $\{(0, t) \in \mathbb{H} \mid |t| \leq \pi/2\}$ is contained in ∂C . But there is no geodesic with curvature $H = 2$ which starts from Σ^- and passes through a point $(0, t)$ with $|t| < \pi/2$. This contradicts Claim 1.

For all $z \in \partial D$, there is a geodesic contained in ∂C , with endpoints $(0, -\pi/2)$, $(0, \pi/2)$, and which passes through $(z, f(z))$. The projection of this geodesic onto the xy -plane is a (full) circle with radius $1/2$ passing through the points 0 and z . Hence ∂C contains the surface of rotation around the t -axis generated by the curve (2.4), and the claim follows. $\square$

References

- [A] Ambrosio, L.: Transport equation and Cauchy problem for BV vector fields. *Invent. Math.* **158** (2004), 227–260
- [AFP] Ambrosio, L., Fusco, N., Pallara, D.: *Functions of Bounded Variation and Free Discontinuity Problems*. Oxford Mathematical Monographs, Oxford Science Publications, Clarendon Press, Oxford, 2000
- [CDST] Capogna, L., Danielli D., Scott D. P., Tyson, J. T.: *An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem*. Preprint, 2006

- [CHY] Cheng, J.-H., Hwang, J.-F., Yang, P.: Existence and uniqueness for p -area minimizers in the Heisenberg group. Preprint, 2006
- [CHMY] Cheng, J.-H., Hwang, J.-F., Malchiodi, A., Yang, P.: Minimal surfaces in pseudohermitian geometry. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **4** (2005), 129–177
- [DGN1] Danielli, D., Garofalo, N., Nhieu, D.-M.: Minimal surfaces, surfaces of constant mean curvature and isoperimetry in Carnot groups. Preprint, 2001
- [DGN2] Danielli, D., Garofalo, N., Nhieu, D.-M.: A partial solution of the isoperimetric problem for the Heisenberg group. Preprint, 2006
- [DL] DiPerna, R. J., Lions, P. L.: Ordinary differential equations, transport theory and Sobolev spaces. *Invent. Math.* **98** (1989), 511–547
- [FGW] Franchi, B., Gallot, S., Wheeden, R. L.: Sobolev and isoperimetric inequalities for degenerate metrics. *Math. Ann.* **300** (1994), 557–571
- [FSSC] Franchi, B., Serapioni, R., Serra Cassano, F.: Rectifiability and perimeter in the Heisenberg group. *Math. Ann.* **321** (2001), 479–531
- [GN] Garofalo, N., Nhieu, D.-M.: Isoperimetric and Sobolev inequalities for Carnot–Carathéodory spaces and the existence of minimal surfaces. *Comm. Pure Appl. Math.* **49** (1996), 1081–1144
- [LM] Leonardi, G. P., Masnou, S.: On the isoperimetric problem in the Heisenberg group $\mathbb{H}^n$. *Ann. Mat. Pura Appl. (4)* **184** (2005), 533–553
- [LR] Leonardi, G. P., Rigot, S.: Isoperimetric sets on Carnot groups. *Houston J. Math.* **29** (2003), 609–637
- [M] Montefalcone, F.: Some relations among volume, intrinsic perimeter and one-dimensional restrictions of BV functions in Carnot groups. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **4** (2005), 79–128
- [Mo] Monti, R.: Brunn–Minkowski and isoperimetric inequality in the Heisenberg group. *Ann. Acad. Sci. Fenn. Math.* **28** (2003), 99–109
- [MM] Monti, R., Morbidelli, D.: Isoperimetric inequality in the Grushin plane. *J. Geom. Anal.* **14** (2004), 355–368
- [MSC] Monti, R., Serra Cassano, F.: Surface measures in Carnot–Carathéodory spaces. *Calc. Var. Partial Differ. Equ.* **13** (2001), 339–376
- [P1] Pansu, P.: Une inégalité isopérimétrique sur le groupe de Heisenberg. *C. R. Acad. Sci. Paris Sér. I Math.* **295** (1982), 127–130
- [P2] Pansu, P.: An isoperimetric inequality on the Heisenberg group. Conference on differential geometry on homogeneous spaces (Turin, 1983). *Rend. Sem. Mat. Univ. Politec. Torino, Special Issue* (1983), 159–174

- [Pa1] Pauls, S. D.: Minimal surfaces in the Heisenberg group. *Geom. Dedicata* **104** (2004), 201–231
- [Pa2] Pauls, S. D.: H-minimal graphs of low regularity in $\mathbb{H}^1$. *Comment. Math. Helv.* **81** (2006), 337–381
- [R] Ritoré, M.: A proof by calibration of an isoperimetric inequality in the Heisenberg group, work in progress
- [RR1] Ritoré, M., Rosales, C.: Rotationally invariant hypersurfaces with constant mean curvature in the Heisenberg group $\mathbb{H}^n$. Preprint, 2005
- [RR2] Ritoré, M., Rosales, C.: Area-stationary surfaces in the Heisenberg group $\mathbb{H}^1$. Preprint, 2005